

# Path Integrals on a Compact Manifold with Non-negative Curvature

Adrian P. C. Lim

Department of Mathematics, Cornell University

Email: al382@cornell.edu

## Abstract

A typical path integral on a manifold,  $M$  is an informal expression of the form

$$\frac{1}{Z} \int_{\sigma \in H(M)} f(\sigma) e^{-E(\sigma)} \mathcal{D}\sigma,$$

where  $H(M)$  is a Hilbert manifold of paths with energy  $E(\sigma) < \infty$ ,  $f$  is a real valued function on  $H(M)$ ,  $\mathcal{D}\sigma$  is a “Lebesgue measure ” and  $Z$  is a normalization constant. For a compact Riemannian manifold  $M$ , we wish to interpret  $\mathcal{D}\sigma$  as a Riemannian “volume form ” over  $H(M)$ , equipped with its natural  $G^1$  metric. Given an equally spaced partition,  $\mathcal{P}$  of  $[0, 1]$ , let  $H_{\mathcal{P}}(M)$  be the finite dimensional Riemannian submanifold of  $H(M)$  consisting of piecewise geodesic paths adapted to  $\mathcal{P}$ . Under certain curvature restrictions on  $M$ , it is shown that

$$\frac{1}{Z_{\mathcal{P}}} e^{-\frac{1}{2}E(\sigma)} dVol_{H_{\mathcal{P}}}(\sigma) \rightarrow \rho(\sigma) d\nu(\sigma) \text{ as } \text{mesh}(\mathcal{P}) \rightarrow 0,$$

where  $Z_{\mathcal{P}}$  is a “normalization” constant,  $E : H(M) \rightarrow [0, \infty)$  is the energy functional,  $Vol_{H_{\mathcal{P}}}$  is the Riemannian volume measure on  $H_{\mathcal{P}}(M)$ ,  $\nu$  is Wiener measure on continuous paths in  $M$ , and  $\rho$  is a certain density determined by the curvature tensor of  $M$ .

## 1 Introduction

Suppose we have a Riemannian manifold  $(M, g)$  of dimension  $d$  with metric  $g$ . We will only consider  $M$  to be compact or  $\mathbb{R}^d$ . Fix a point  $o$  on the manifold  $M$  and let  $V : M \rightarrow \mathbb{R}$  be a potential function. In classical mechanics, the path  $\sigma : [0, T] \rightarrow M$ ,  $\sigma(0) = o$ , subject to the potential  $V$ , can be obtained by solving Newton’s equation of motion

$$\frac{\nabla}{dt} \sigma'(t) = -\text{grad } V(\sigma(t)),$$

where given any vector field  $X(s)$  on  $\sigma(s)$ , define

$$\frac{\nabla X(s)}{ds} := //_s(\sigma) \frac{d}{ds} \{ //_s^{-1}(\sigma) X(s) \}, \quad (1.1)$$

and  $//_s(\sigma) : T_o M \rightarrow T_{\sigma(s)} M$  is parallel translation along  $\sigma$  relative to the Levi Civita covariant derivative  $\nabla$ . The Hamiltonian of the system,  $H$  is then given by

$$H(\sigma(t), \sigma'(t)) = \frac{1}{2} \| \sigma'(t) \|^2 + V(\sigma(t)),$$

where  $\| v \|^2 := g(v, v)$  and mass is set to be 1.

In Quantum Mechanics, observables are no longer functions, but rather Hermitian operators on some Hilbert space. Let  $q = (q_1, q_2, \dots, q_d)$  be the cartesian coordinates on  $\mathbb{R}^d$  and  $p_i$  be the momentum corresponding to  $q_i$ . In canonical quantization on  $\mathbb{R}^d$ , the quantum mechanical operator  $\hat{H}$  corresponding to the classical Hamiltonian,  $H = \frac{1}{2} \sum_i p_i^2 + V(q)$ , is given by

$$\hat{H} = -\frac{\hbar^2}{2} \sum_i \frac{\partial^2}{\partial q_i^2} + M_V$$

where  $\hbar = \frac{h}{2\pi}$ ,  $h$  is Planck's constant and  $M_V$  is multiplication operator by  $V$ .

However, on a manifold, one aims to quantize the Hamiltonian  $H = \frac{1}{2} g^{ij}(q) p_i p_j + V(q)$  where  $q$  is some coordinate system on  $M$ . Using the ‘‘Feynman's (Kac) path integral prescription’’, one defines the operator  $e^{-T\hat{H}}$  via an integral

$$(e^{-T\hat{H}} f)(o) := \frac{1}{Z_T} \int_{H_T(M)} e^{-\int_0^T H(\sigma(t), \sigma'(t)) dt} f(\sigma(T)) \mathcal{D}\sigma, \quad (1.2)$$

where  $H(\sigma(t), \sigma'(t)) = \frac{1}{2} \| \sigma'(t) \|^2 + V(\sigma(t))$  is the classical Hamiltonian.  $H_T(M)$  is the space of finite energy paths,  $Z_T$  is some normalization constant and  $\mathcal{D}\sigma$  is to be interpreted as a ‘‘Lebesgue type measure’’. The operator  $\hat{H}$  can then be obtained by differentiating the operator  $e^{-T\hat{H}}$  with respect to  $T$  at 0.

For simplicity, we will assume that  $T = 1$  and set  $\hbar = 1$ . Furthermore we will set  $V = 0$  since quantizing a scalar function on the manifold  $M$  corresponds to the multiplication operator with  $V$ .

**Definition 1.1** Define  $H(M)$ , a Hilbert manifold of absolutely continuous paths with finite energy,

$$H(M) = \{ \sigma : [0, 1] \rightarrow M \mid \sigma(0) = o \in M \text{ and } E(\sigma) < \infty \} \quad (1.3)$$

where the energy  $E$  is given by

$$E(\sigma) := \int_0^1 g(\sigma'(s), \sigma'(s)) ds. \quad (1.4)$$

The tangent space  $T_\sigma H(M)$  to  $H(M)$  at  $\sigma$  may be identified with the space of absolutely continuous vector fields along  $\sigma$ . On this Hilbert manifold  $H(M)$ , we can define a metric  $G^1$  as follows.

**Definition 1.2** *Let  $T_\sigma H(M)$  be the space of absolutely continuous vector fields  $X$  along  $\sigma$  (i.e.  $X(s) \in T_{\sigma(s)} M \forall s \in [0, 1]$ ) such that  $G^1(X, X) < \infty$  where*

$$G^1(X, X) := \int_0^1 g \left( \frac{\nabla X(s)}{ds}, \frac{\nabla X(s)}{ds} \right) ds, \quad (1.5)$$

and  $\frac{\nabla}{ds}$  is defined as in Equation (1.1).

See [Eel58, Eli67, FK72, Kli78, Pal63] for more details. By polarization, Equation (1.5) defines a Riemannian metric on  $H(M)$ .

The integral over  $H(M)$ , defined in Equation (1.2) however, is highly heuristic. Firstly, the normalization constant  $Z_1$  may be interpreted to be 0 or  $\infty$ . Secondly,  $\mathcal{D}\sigma$  which is to be interpreted as “Lebesgue measure”, fails to exist in an infinite dimensional space.

We would like to make sense out of the RHS of Equation (1.2), by writing it as a limit of a sequence of integrals over finite dimensional spaces  $H_{\mathcal{P}}(M)$ .

**Definition 1.3** *Let*

$$\mathcal{P} = \{0 = s_0 < s_1 < s_2 < \dots < s_n = 1\} \quad (1.6)$$

*be a partition of  $[0, 1]$ . Define  $H_{\mathcal{P}}(M)$  as a set of piecewise geodesics paths in  $H(M)$  which change directions only at the partition points.*

$$H_{\mathcal{P}}(M) = \left\{ \sigma \in H(M) \cap C^2(I \setminus \mathcal{P}) \mid \frac{\nabla \sigma'(s)}{ds} = 0 \text{ for } s \notin \mathcal{P} \right\}.$$

It will be shown later that  $H_{\mathcal{P}}(M)$  is a finite dimensional submanifold of  $H(M)$ . In fact,  $H_{\mathcal{P}}(M)$  is diffeomorphic to  $(\mathbb{R}^d)^n$ . For  $\sigma \in H_{\mathcal{P}}(M)$ , the tangent space  $T_\sigma H_{\mathcal{P}}(M)$  can be identified with elements  $X \in T_\sigma H(M)$  satisfying the Jacobi equations on  $I \setminus \mathcal{P}$ . As a submanifold of  $H(M)$ ,  $H_{\mathcal{P}}(M)$  inherits the induced metric  $G^1|_{TH_{\mathcal{P}}(M)}$  by restricting the  $G^1$  metric on  $H(M)$ .

If  $N^p$  is any manifold with a metric  $G$ , define a volume density  $Vol_G$  on  $T_n N$  by

$$Vol_G(v_1, v_2, \dots, v_p) = \sqrt{\det \{G(v_i, v_j)\}_{i,j=1}^p} \quad (1.7)$$

where  $\{v_1, v_2, \dots, v_p\} \subset T_n N$  is a basis and  $n \in N$ .

**Theorem 1.4** *Given a density of the form  $\rho \text{Vol}_G$ , where  $\rho : N \rightarrow [0, \infty)$ , there exists a unique measure  $m_G$  on  $N$  such that*

$$\int_{\mathcal{D}(y)} f \, dm_G = \int_{\mathcal{D}(y)} f \rho \, \text{Vol}_G \left( \frac{\partial}{\partial y_1}, \frac{\partial}{\partial y_2}, \dots, \frac{\partial}{\partial y_p} \right) dy_1 \dots dy_p$$

*for any local coordinates  $y = (y_1, \dots, y_p) : \mathcal{D}(y) \rightarrow \mathbb{R}^p$  and measurable function  $f : N \rightarrow [0, \infty)$ . If  $\rho = 1$ , the associated measure will be called Riemann volume measure.*

**Definition 1.5** *Let  $\text{Vol}_{\mathcal{P}}$  denote the density on  $H_{\mathcal{P}}(M)$  determined by  $G^1|_{TH_{\mathcal{P}}(M) \otimes TH_{\mathcal{P}}(M)}$  using Equation (1.7).*

Given the above definition, we can now define a measure on  $H_{\mathcal{P}}(M)$ .

**Definition 1.6** *For each partition  $\mathcal{P}$  of  $[0, 1]$  as in Equation (1.6), let  $\nu_{\mathcal{P}}$  denote the unique measure on  $H_{\mathcal{P}}(M)$  as in Theorem 1.4, defined by the density*

$$\frac{1}{Z_{\mathcal{P}}^1} e^{-\frac{1}{2}E} \text{Vol}_{\mathcal{P}}$$

*where  $E : H(M) \rightarrow [0, \infty)$  is the energy functional defined in Equation (1.4) and  $Z_{\mathcal{P}}^1$  is a normalization constant given by*

$$Z_{\mathcal{P}}^1 = (2\pi)^{\frac{dn}{2}}. \quad (1.8)$$

We can now write the RHS of Equation (1.2) as a limit of a sequence of integrals over the finite dimensional space  $H_{\mathcal{P}}(M)$ , equipped with the measure  $\nu_{\mathcal{P}}$ . Our result shows that this limit can be written as an integral over the Wiener space of  $M$ , with Wiener measure  $\nu$ . (See Definition 1.8 below.)

**Definition 1.7** *Define  $\Delta_i s = s_i - s_{i-1}$  and  $|\mathcal{P}| = \bigvee_{i=1, \dots, n} \Delta_i s = \max\{\Delta_i s : i = 1, 2, \dots, n\}$  be the norm of the partition and  $J_i := (s_{i-1}, s_i]$  for  $i = 1, 2, \dots, n$ .*

Let  $\Delta = \text{tr } \nabla^2$  denote the Laplacian acting on  $C^\infty(M)$  and  $p_s(x, y)$  be the fundamental solution to the heat equation,

$$\frac{\partial u}{\partial s} = \frac{1}{2} \Delta u.$$

In the case when  $M = \mathbb{R}^d$ ,

$$p_s(x, y) = \left( \frac{1}{2\pi s} \right)^{\frac{d}{2}} e^{-\frac{1}{2s} \|x-y\|^2}.$$

**Definition 1.8** *The Wiener space  $W(M)$  is the path space*

$$W(M) = \{\sigma : [0, 1] \rightarrow M : \sigma(0) = o \text{ and } \sigma \text{ is continuous}\}.$$

*The Wiener measure  $\nu$  associated to  $(M, g, o)$  is the unique probability measure on  $W(M)$  such that*

$$\int_{W(M)} f(\sigma) d\nu(\sigma) = \int_{M^n} F(x_1, \dots, x_n) \prod_{i=1}^n p_{\Delta_i s}(x_{i-1}, x_i) dm(x_1) \cdots dm(x_n) \quad (1.9)$$

*for all functions  $f$  of the form  $f(\sigma) = F(\sigma(s_1), \dots, \sigma(s_n))$ , for all  $\mathcal{P}$ , a partition of  $[0, 1]$  as in Equation (1.6) and  $F : M^n \rightarrow \mathbb{R}$  is a bounded measurable function. In Equation (1.9),  $dm(x)$  denotes the Riemann volume measure on  $M$  as in Theorem 1.4 and by convention  $x_0 := o$ .*

It is known that there exists a unique probability measure  $\nu$  on  $W(M)$  satisfying Equation (1.9). The measure  $\nu$  is concentrated on continuous but nowhere differentiable paths.

**Notation 1.9** *When  $M = \mathbb{R}^d$ ,  $g(\cdot, \cdot)$  is the usual dot product and  $o = 0$ , the measure  $\nu$  defined in Definition 1.8 is the standard Wiener measure on  $W(\mathbb{R}^d)$ . We will denote this standard Wiener measure by  $\mu$  rather than  $\nu$ . We will also let  $b(s) : W(\mathbb{R}^d) \rightarrow \mathbb{R}^d$  be the coordinate map such that*

$$b(s)(\omega) := \omega(s)$$

*for all  $\omega \in W(\mathbb{R}^d)$ .*

**Remark 1.10** *The process  $\{b(s)\}_{s \in [0, 1]}$  is a standard  $\mathbb{R}^d$ -valued Brownian motion on the probability space  $(W(\mathbb{R}^d), \mu)$ .*

Suppose we now view  $M$  as an imbedded submanifold of  $\mathbb{R}^d$  with the induced Riemannian structure. Let  $P(m)$  be the projection on the tangent space  $T_m M$  and  $v \in T_m M$ . Then for a vector field  $X(m)$ ,

$$\nabla_v X = P(\sigma(0)) \frac{d}{dt} X(\sigma(t)) \Big|_{t=0}$$

where  $\sigma$  is a path in  $M$  such that  $\sigma(0) = m$  and  $\sigma'(0) = v$ . Define a projection  $Q$  on the orthogonal complement of  $T_m M$  by  $Q = I - P$ , where  $I$  is the identity. With this definition, for any vector  $v \in T_m M$ , one can define parallel translation along  $\sigma$  by  $//_s(\sigma)v := w(s)v$  where  $w$  solves the following differential equation

$$w'(t) + dQ(\sigma'(t))w(t) = 0, \quad w(0) = P(\sigma(0)).$$

**Theorem 1.11** *Let  $\Sigma$  be an  $M$ -valued semi-martingale and  $V_0(m)$  be a measurable vector field on  $M$ , then there is a unique parallel  $TM$ -valued semi-martingale  $V$  such that  $V_0 = V_0(\Sigma_0)$  and  $V_s \in T_{\Sigma_s}M$  for all  $s$ . More explicitly,  $V_s = w_s V(\Sigma_0)$  where  $w$  solves the following Stratonovich stochastic differential equation*

$$\delta w + dQ(\delta \Sigma)w = 0, \quad w_0 = P(\Sigma_0). \quad (1.10)$$

For a proof of this theorem, the reader should refer to [Rao04]. Thus we can now define a “stochastic” extension of parallel translation.

**Definition 1.12** (Stochastic Parallel Translation) *Given  $v \in T_{\Sigma_0}M$  and  $M$ -valued semi-martingale  $\Sigma$ , define stochastic parallel translation  $\widetilde{\parallel}$  by*

$$\widetilde{\parallel}_s v := w_s v$$

where  $w$  solves Equation (1.10). This is going to be used for the particular semi-martingale  $\Sigma_s(\sigma) := \sigma(s)$  on  $(W(M), \nu)$ .

**Definition 1.13** *The curvature tensor  $R$  of  $\nabla$  is*

$$R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]}Z$$

for all vector fields  $X, Y$  and  $Z$  on  $M$ . The sectional curvature  $S(V)$  where  $V \subseteq T_m M$  with  $\dim(V) = 2$ , is defined by

$$S(V) = \frac{g(R(u, v)u, v)}{\|u\|^2 \|v\|^2 - g(u, v)^2},$$

where  $\{u, v\}$  is a basis for  $V$ . It can be shown that this definition is independent of the basis used. Let  $\{e_i\}_{i=1}^d \subseteq T_m M$  be an orthonormal frame at  $m \in M$ . The Ricci tensor of  $(M, g)$  is  $\text{Ric } v = \sum_{i=1}^d R(v, e_i)e_i$ , the scalar curvature  $\text{Scal}$  is  $\text{Scal} = \sum_{i=1}^d g(\text{Ric } e_i, e_i)$  and  $\Gamma_m$  is given by

$$\Gamma_m = \sum_{i,j=1}^d \left( R(e_i, R(e_i, \cdot)e_j)e_j + R(e_i, R(e_j, \cdot)e_i)e_j + R(e_i, R(e_j, \cdot)e_j)e_i \right).$$

Define for any  $\sigma \in W(M)$ ,  $K_\sigma : L^2([0, 1] \rightarrow T_o M) \longrightarrow L^2([0, 1] \rightarrow T_o M)$  by

$$(K_\sigma v)(s) = \int_0^1 (s \wedge t) \widetilde{\parallel}_t^{-1}(\sigma) \left( \Gamma_{\sigma(t)} \widetilde{\parallel}_t(\sigma) v(t) \right) dt$$

where  $\widetilde{\parallel}$  is stochastic parallel translation.

It will be proved later that  $K_\sigma$  is a trace class operator. We can now state the main result.

**Theorem 1.14** *Let  $(M, g)$  be a compact Riemannian manifold with dimension  $d$ . Let  $\mathcal{P} = \{0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}, 1\}$  be an equally spaced partition. Suppose that  $f : W(M) \rightarrow \mathbb{R}$  is bounded and continuous and that  $0 \leq S < \frac{3}{17d}$ , then*

$$\begin{aligned} & \lim_{|\mathcal{P}| \rightarrow 0} \int_{H_{\mathcal{P}}(M)} f(\sigma) d\nu_{\mathcal{P}}(\sigma) \\ &= \int_{W(M)} f(\sigma) e^{-\frac{1}{6} \int_0^1 \text{Scal}(\sigma(s)) ds} \sqrt{\det \left( I + \frac{1}{12} K_{\sigma} \right)} d\nu(\sigma). \end{aligned} \quad (1.11)$$

## 1.1 Known Results

Using  $H_{\mathcal{P}}(M)$  to approximate the Wiener space  $W(M)$  was done in [AD99]. However, the choice of metrics used on  $TH_{\mathcal{P}}(M)$  in [AD99] are different from  $G^1|_{TH_{\mathcal{P}}(M)}$ .

**Definition 1.15** *For each partition  $\mathcal{P}$  of  $[0, 1]$  as in Equation (1.6), let  $G_{\mathcal{P}}^1, G_{\mathcal{P}}^0$  be the metrics on  $TH_{\mathcal{P}}(M)$ , given by*

$$\begin{aligned} G_{\mathcal{P}}^1(X, Y) &= \sum_{i=1}^n g \left( \frac{\nabla X(s_{i-1}+)}{ds}, \frac{\nabla Y(s_{i-1}+)}{ds} \right) \Delta_i s, \\ G_{\mathcal{P}}^0(X, Y) &= \sum_{i=1}^n g ( X(s_{i-1}+), Y(s_{i-1}+) ) \Delta_i s. \end{aligned}$$

for all  $X, Y \in T_{\sigma}H_{\mathcal{P}}(M)$  and  $\sigma \in H_{\mathcal{P}}(M)$ . Note that  $\frac{\nabla X(s_{i-1}+)}{ds} = \lim_{s \rightarrow s_{i-1}+} \frac{\nabla X(s)}{ds}$ .

Observe that  $G_{\mathcal{P}}^1$  is some sort of Riemann sum approximation to  $G^1$ .

**Definition 1.16** *For each partition  $\mathcal{P}$  of  $[0, 1]$  as in Equation (1.6), define unique measures  $\nu_{G_{\mathcal{P}}^1}$  and  $\nu_{G_{\mathcal{P}}^0}$  on  $H_{\mathcal{P}}(M)$ , as in Theorem 1.4, defined by densities*

$$\frac{1}{Z_{\mathcal{P}}^1} e^{-\frac{1}{2}E} \text{Vol}_{G_{\mathcal{P}}^1} \text{ and } \frac{1}{Z_{\mathcal{P}}^0} e^{-\frac{1}{2}E} \text{Vol}_{G_{\mathcal{P}}^0}$$

respectively, where  $E : H(M) \rightarrow [0, \infty)$  is the energy functional defined in Equation (1.4) and  $\text{Vol}_{G_{\mathcal{P}}^1}$  and  $\text{Vol}_{G_{\mathcal{P}}^0}$  are determined by  $G_{\mathcal{P}}^1$  and  $G_{\mathcal{P}}^0$  respectively using Equation (1.7).  $Z_{\mathcal{P}}^1$  is a normalization constant given by Equation (1.8) and

$$Z_{\mathcal{P}}^0 = \prod_{i=1}^n \left( \sqrt{2\pi} (s_i - s_{i-1}) \right)^d.$$

The following theorem was proved in [AD99].

**Theorem 1.17** *Let  $M$  be a compact Riemannian manifold. Suppose that  $f : W(M) \rightarrow \mathbb{R}$  is bounded and continuous, then*

$$\begin{aligned} \lim_{|\mathcal{P}| \rightarrow 0} \int_{H_{\mathcal{P}}(M)} f(\sigma) d\nu_{G_{\mathcal{P}}^1}(\sigma) &= \int_{W(M)} f(\sigma) d\nu(\sigma) \text{ and} \\ \lim_{|\mathcal{P}| \rightarrow 0} \int_{H_{\mathcal{P}}(M)} f(\sigma) d\nu_{G_{\mathcal{P}}^0}(\sigma) &= \int_{W(M)} f(\sigma) e^{-\frac{1}{6} \int_0^1 \text{Scal}(\sigma(s)) ds} d\nu(\sigma). \end{aligned}$$

Using the Feynman-Kac formula,  $\hat{H}$  in Equation (1.2) is given by

$$\hat{H} = -\frac{1}{2}\Delta + \kappa \text{Scal}$$

where  $\Delta$  is the Laplacian operator and  $\kappa = 0, \frac{1}{6}$  for the  $G_{\mathcal{P}}^1$  and  $G_{\mathcal{P}}^0$  metrics respectively. However, the integral in Equation (1.11) is not of the Feynman-Kac form. Hence the interpretation of the operator  $\hat{H}$  corresponding to this integral is unclear at this stage.

## 2 Finite Dimensional Approximations

A detailed account of this section is given in [AD99].

Let  $\pi : O(M) \rightarrow M$  denote the bundle of orthogonal frames on  $M$ . An element  $u \in O(M)$  is an isometry

$$u : \mathbb{R}^d \rightarrow T_{\pi(u)}M.$$

Fix  $u_o \in \pi^{-1}(o)$ , which identifies  $T_oM$  of  $M$  at  $o$  with  $\mathbb{R}^d$ .

Define  $\theta$ , a  $\mathbb{R}^d$ -valued form on  $O(M)$  by  $\theta_u(\xi) = u^{-1}\pi_*\xi$  for all  $u \in O(M)$ ,  $\xi \in T_uO(M)$  and let  $\vartheta$  be the  $so(d)$ -valued connection form on  $O(M)$  defined by  $\nabla$ . Explicitly, if  $s \rightarrow u(s)$  is a smooth path in  $O(M)$  then

$$\vartheta(u'(0)) = u(0)^{-1} \frac{\nabla u(s)}{ds} \Big|_{s=0}$$

where  $\frac{\nabla}{ds}$  is defined as in Equation (1.1) with  $X$  replaced by  $u$ . We define the horizontal lift  $\mathcal{H}_u : T_{\pi(u)}M \rightarrow T_uO(M)$  by

$$\theta \mathcal{H}_u u = id_{\mathbb{R}^d}, \quad \vartheta_u \mathcal{H}_u = 0.$$

Explicitly, for  $v \in T_{\pi(u)}M$ ,  $\mathcal{H}_u v = \frac{d}{dt}|_{t=0} / t(\sigma)u$  where  $\dot{\sigma}(0) = v$ .

**Definition 2.1** *For  $a, c \in \mathbb{R}^d$ , let*

$$\Omega_u(a, c) := u^{-1}R(ua, uc)u.$$



Let  $H(O(M))$  be the set of finite energy paths  $u : [0, 1] \rightarrow O(M)$  as defined in Equation (1.3) with  $M$  replaced by  $O(M)$  and  $o$  by  $u_o$ . For  $\sigma \in H(M)$ , let  $u$  be defined by the ordinary differential equation

$$u'(s) = \mathcal{H}_{u(s)}\sigma'(s), \quad u(0) = u_o.$$

This equation implies that  $\vartheta(u'(s)) = 0$  or that  $\frac{\nabla u(s)}{ds} = 0$ . Thus we have

$$u(s) = //_s(\sigma)u_o$$

where  $//_s(\sigma)$  is parallel translation along  $\sigma$ . Since  $u_o$  is fixed, we will drop it and write  $u(s) = //_s$ . We will call  $u(s)$  a horizontal lift of  $\sigma$  starting at  $u_o$  and use it to define  $\phi$ , which associates  $\omega \in H(\mathbb{R}^d)$  with a path  $\sigma \in H(M)$ .

**Definition 2.2** (Cartan's Development Map) *The development map,  $\phi : H(\mathbb{R}^d) \rightarrow H(M)$  is defined for  $\omega \in H(\mathbb{R}^d)$  by  $\phi(\omega) = \sigma \in H(M)$  where  $\sigma$  solves the functional differential equation*

$$\sigma'(s) = //_s(\sigma)\omega'(s), \quad \sigma(0) = o. \quad (2.1)$$

The development map,  $\phi$  is smooth and injective. We can define an anti-development map,  $\phi^{-1} : H(M) \rightarrow H(\mathbb{R}^d)$  given by  $\omega = \phi^{-1}(\sigma)$  where

$$\omega(s) = \int_0^s //_r^{-1}(\sigma)\sigma'(r) \, dr.$$

Again,  $\phi^{-1}$  is smooth and is injective. Therefore,  $\phi : H(\mathbb{R}^d) \rightarrow H(M)$  is a diffeomorphism of infinite dimensional Hilbert manifolds.

**Definition 2.3** *For each  $h \in C^\infty(H(M) \rightarrow H(\mathbb{R}^d))$  and  $\sigma \in H(M)$ , let  $X^h(\sigma) \in T_\sigma H(M)$  be given by*

$$X_s^h(\sigma) := //_s(\sigma)h_s(\sigma) \quad (2.2)$$

for all  $s \in [0, 1]$ , where we have written  $h_s(\sigma)$  as  $h(\sigma)(s)$ .

Define  $H_{\mathcal{P}}(\mathbb{R}^d) = \{\omega \in H \cap C^2(I \setminus \mathcal{P}) \mid \omega''(s) = 0 \text{ for } s \notin \mathcal{P}\}$ , the set of piecewise linear paths in  $H(\mathbb{R}^d)$ , which changes directions only at the partition points.

**Remark 2.4** *The development map  $\phi : H(\mathbb{R}^d) \rightarrow H(M)$  has the property that  $\phi(H_{\mathcal{P}}(\mathbb{R}^d)) = H_{\mathcal{P}}(M)$  where  $H_{\mathcal{P}}(M)$  has been defined in Definition 1.3. If  $\sigma = \phi(\omega)$  with  $\omega \in H_{\mathcal{P}}(\mathbb{R}^d)$ , then differentiating Equation (2.1) gives*

$$\frac{\nabla \sigma'(s)}{ds} = \frac{\nabla}{ds} \left( //_s(\sigma)\omega'(s) \right) = //_s(\sigma)\omega''(s) = 0 \text{ for all } s \notin \mathcal{P}.$$

Because  $\phi : H(\mathbb{R}^d) \rightarrow H(M)$  is a diffeomorphism and  $H_{\mathcal{P}}(\mathbb{R}^d) \subset H(\mathbb{R}^d)$  is an embedded submanifold, so it follows that  $H_{\mathcal{P}}(M)$  is an embedded submanifold of  $H(M)$ . Therefore for each  $\sigma \in H_{\mathcal{P}}(M)$ ,  $T_{\sigma}H_{\mathcal{P}}(M)$  may be viewed as a subspace of  $T_{\sigma}H(M)$ . The next proposition identifies this subspace. See [AD99] for a proof.

**Proposition 2.5** *Let  $\sigma \in H_{\mathcal{P}}(M)$ , then  $X \in T_{\sigma}H(M)$  is in  $T_{\sigma}H_{\mathcal{P}}(M)$  if and only if*

$$\frac{\nabla^2}{ds^2}X(s) = R(\sigma'(s), X(s))\sigma'(s) \quad (2.3)$$

*on  $I \setminus \mathcal{P}$ . Equivalently, letting  $\omega = \phi^{-1}(\sigma)$ ,  $u = //(\sigma)$  and  $h \in H(\mathbb{R}^d)$ , then  $X^h \in T_{\sigma}H(M)$  defined in Equation (2.2) is in  $T_{\sigma}H_{\mathcal{P}}(M)$  if and only if*

$$h''(s) = \Omega_{u(s)}(\omega'(s), h(s))\omega'(s) \quad (2.4)$$

*on  $I \setminus \mathcal{P}$ .*

## 2.1 Comparing $\nu_{\mathcal{P}}$ and $\nu_{G_{\mathcal{P}}^1}$ on $M$

**Definition 2.6** *For  $\omega \in H_{\mathcal{P}}(\mathbb{R}^d)$ , let  $\{h_{k,a}\}_{\substack{k=1,2,\dots,n \\ a=1,2,\dots,d}}$  be any basis in  $\phi_*^{-1}(T_{\phi(\omega)}H_{\mathcal{P}}(M)) = H_{\mathcal{P}}(T_{\omega}M)$  and  $\sigma = \phi(\omega)$ . Let  $Vol_{\mathcal{P}}$  be the density associated to  $G^1|_{T_{H_{\mathcal{P}}}(M) \otimes T_{H_{\mathcal{P}}}(M)}$  metric and  $Vol_{G_{\mathcal{P}}^1}$  be the density associated to  $G_{\mathcal{P}}^1$  metric. Then*

$$X_s^{h_{k,a}}(\sigma) := //_s(\sigma) h_{k,a}(s) \text{ for } k = 1, 2, \dots, n \text{ and } a = 1, 2, \dots, d$$

*is a basis for  $T_{\sigma}H_{\mathcal{P}}(M)$  and we define*

$$\begin{aligned} \rho_{\mathcal{P}} &= \frac{|Vol_{\mathcal{P}}(\{X^{h_{k,a}}\}_{\substack{i=1,2,\dots,n \\ a=1,2,\dots,d}})|}{|Vol_{G_{\mathcal{P}}^1}(\{X^{h_{k,a}}\}_{\substack{i=1,2,\dots,n \\ a=1,2,\dots,d}})|} \\ &= \frac{\sqrt{\det(\{G^1(X^{h_{k,a}}, X^{h_{k',a'}})\}_{(k,a),(k',a')})}}{\sqrt{\det(\{G_{\mathcal{P}}^1(X^{h_{k,a}}, X^{h_{k',a'}})\}_{(k,a),(k',a')})}}. \end{aligned} \quad (2.5)$$

The relevance of this definition is contained in the next remark.

**Remark 2.7** *First off, it is well known (and easily verified) that the  $\rho_{\mathcal{P}}(\sigma)$  defined in Equation (2.5) is well defined independent of the choice of basis  $\{h_{k,a}\}_{\substack{k=1,2,\dots,n \\ a=1,2,\dots,d}}$ . Secondly, if  $\nu_{\mathcal{P}}$  and  $\nu_{G_{\mathcal{P}}^1}$  are the measures associated to  $G^1|_{T_{H_{\mathcal{P}}}(M) \otimes T_{H_{\mathcal{P}}}(M)}$  and  $G_{\mathcal{P}}^1$  respectively, then  $d\nu_{\mathcal{P}} = \rho_{\mathcal{P}} \cdot d\nu_{G_{\mathcal{P}}^1}$ .*

From [AD99], we know the limiting behavior of the measure  $\nu_{G_{\mathcal{P}}^1}$ . Hence, our proof that  $\nu_{\mathcal{P}}$  has a limit will break into two parts. Very roughly speaking we are going to first show that  $\{\rho_{\mathcal{P}} : \mathcal{P}\}$  is uniformly integrable and then we will show that  $\lim_{|\mathcal{P}| \rightarrow 0} \rho_{\mathcal{P}}$  exists in  $\mu$ -measure. This rough outline will have to be appropriately modified since  $\{\rho_{\mathcal{P}} : \mathcal{P}\}$  are functions on different probability spaces for each  $\mathcal{P}$ . This will be remedied by pulling  $\rho_{\mathcal{P}}$  to classical Wiener space  $(W(T_o M), \mu)$  using Cartan's rolling map  $\phi$  and the natural piecewise linear approximation map from  $W(T_o M)$  to  $H_{\mathcal{P}}(T_o M)$ . We will identify  $T_o(M)$  with  $\mathbb{R}^d$ .

Before we move on, we would like to point out that  $\rho_{\mathcal{P}} \circ \phi$  is only defined on  $H_{\mathcal{P}}(\mathbb{R}^d)$ , which has  $\mu$ -measure zero.

**Definition 2.8** Let  $\{b(s)\}_{s \in [0,1]}$  be the standard  $\mathbb{R}^d$  Brownian motion on  $(W(\mathbb{R}^d), \mu)$  and  $\mathcal{P}$  be any partition, i.e.  $b(s) : W(\mathbb{R}^d) \rightarrow \mathbb{R}^d$ ,

$$b(s)(\omega) := b(s, \omega) := \omega(s).$$

By abuse of notation, define  $b_{\mathcal{P}} : W(\mathbb{R}^d) \rightarrow H_{\mathcal{P}}(\mathbb{R}^d)$  by

$$b_{\mathcal{P}}(s) = b(s_{i-1}) + (s - s_{i-1}) \frac{\Delta_i b}{\Delta_i s} \quad \text{if } s \in (s_{i-1}, s_i]$$

where  $\Delta_i b := b(s_i) - b(s_{i-1})$ . We will write  $b_n = b_{\mathcal{P}_n}$  if  $\mathcal{P}_n = \{0 < \frac{1}{n} < \dots < \frac{n}{n} = 1\}$  is an equally spaced partition

Thus by composing with  $b_{\mathcal{P}}$ , we can now view  $\rho_{\mathcal{P}} \circ \phi \circ b_{\mathcal{P}}$  as a random variable on  $(W(\mathbb{R}^d), \mu)$ .

### 3 Uniform Integrability of $\{\rho_n \circ \phi \circ b_n\}_{n=1}^{\infty}$

We will first show that  $\rho_{\mathcal{P}} \circ \phi \circ b_{\mathcal{P}}$  is uniformly integrable. But first, we need to write down a formula for  $\rho_{\mathcal{P}}$ .

#### 3.1 A First Formula for $\rho_n$

We will now only consider equally spaced partitions  $\mathcal{P}_n = \{0 = s_0 < s_1 < s_2 < \dots < s_n = 1\}$ , such that  $\Delta_i s = \frac{1}{n}$ ,  $i = 1, \dots, n$  and write  $\rho_n = \rho_{\mathcal{P}_n}$ . Let  $\sigma \in H_{\mathcal{P}_n}(M)$  and consider  $\omega = \phi^{-1}(\sigma)$ . On each  $J_i = (s_{i-1}, s_i]$ ,  $i = 1, 2, \dots, n$ ,  $\omega'_i := \omega'(s_{i-1}+)$  is a constant. Thus  $\Delta_i \omega := \omega'_i \Delta_i s$ .

From Proposition 2.5, we know that for  $s \in J_i$ , for each  $h$  such that  $X^h \in TH_{\mathcal{P}_n}(M)$ ,  $h(\omega, s)$  satisfies the ordinary differential equation

$$\frac{d^2 h(\omega, s)}{ds^2} = \Omega_{u(s)}(\omega'_i, h(\omega, s)) \omega'_i. \quad (3.1)$$

Let  $\{e_a\}_{a=1}^d$  be the standard basis for  $\mathbb{R}^d$  and for  $i = 1, 2, \dots, n$ , let

$$e_{i,a} = \left(0, \dots, 0, \overset{i^{\text{th}}}{e_a^{\text{spot}}}, 0, \dots, 0\right) \in (\mathbb{R}^d)^n = \mathbb{R}^{nd}.$$

Then  $\{e_{i,a}\}_{\substack{i=1,\dots,n \\ a=1,\dots,d}}$  is an indexing of the standard basis for  $\mathbb{R}^{nd}$  such that all the components of  $e_{i,a}$  are 0 except at the  $a + (i-1)d$  position, which is 1.

**Notation 3.1** Let  $h_{i,a}(\omega, s)$  denote the continuous function in  $\mathbb{R}^d$  which solves Equation (3.1) on  $[0, 1] \setminus \mathcal{P}_n$  and satisfies

$$h_{i,a}(\omega, 0) = 0, \text{ and } h'_{i,a}(\omega, s_{j-1}+) = \delta_{ij}e_a \text{ for } j = 1, \dots, n.$$

It is easily seen that  $\{h_{i,a}\}_{\substack{i=1,\dots,n \\ a=1,\dots,d}}$  forms a basis for  $\phi_*^{-1}(T_{\phi(\omega)}H_{\mathcal{P}_n}(M)) = TH_{\mathcal{P}_n}(\mathbb{R}^d) \cong H_{\mathcal{P}_n}(\mathbb{R}^d)$ . Further let  $\mathcal{Q}^n$  denote the  $nd \times nd$  matrix which is given in  $d \times d$  blocks,  $\mathcal{Q}^n := \{(\mathcal{Q}_{mk}^n)\}_{m,k=1}^n$ , with

$$(\mathcal{Q}_{mk}^n e_a, e_c)(\omega) := \int_0^1 \langle h'_{ma}(\omega, s), h'_{kc}(\omega, s) \rangle ds \text{ for } a, c = 1, 2, \dots, d.$$

**Notation 3.2** Unless stated otherwise, upper case letters without a superscript  $n$  will denote  $d \times d$  matrices. Upper case letters and scripted upper case letters with a superscript  $n$  will denote  $n \times n$  block matrices with entries being  $d \times d$  blocks. We will reserve  $\mathcal{I}^n$  and  $I$  for a  $nd \times nd$  identity matrix and a  $d \times d$  identity matrix respectively. Matrices with a superscript  $T$  will denote the matrix transpose. To avoid confusion, we will use  $\text{Tr}$  and  $\text{tr}$  to denote taking the trace of a  $nd \times nd$  matrix and a  $d \times d$  matrix respectively.

$b$  will denote a standard  $d$ -dimensional Brownian path in  $\mathbb{R}^d$ . For a piecewise continuous function on  $[0, 1]$ , we will use the notation  $f(s+) = \lim_{r \downarrow s} f(r)$ . We also let  $\langle \cdot, \cdot \rangle$  denote  $g(\cdot, \cdot)|_o$  at base point  $o$ .

**Remark 3.3** All norms used for matrices will be the operator norm. Norms used for vectors will be the euclidean norm.

**Lemma 3.4** The relationship between  $\rho_n \circ \phi$  and  $\mathcal{Q}^n$  is

$$\rho_n \circ \phi = \sqrt{\det(n\mathcal{Q}^n)}. \quad (3.2)$$

**Proof.** Observe that

$$\begin{aligned} G^1(X^{h_{k,a}}, X^{h_{m,c}})(\omega) &= \int_0^1 g\left(\frac{\nabla X^{h_{k,a}}(\omega, s)}{ds}, \frac{\nabla X^{h_{m,c}}(\omega, s)}{ds}\right) ds \\ &= \int_0^1 \langle h'_{k,a}(\omega, s), h'_{m,c}(\omega, s) \rangle ds \\ &= \langle \mathcal{Q}_{mk}^n(\omega) e_a, e_c \rangle = \langle \mathcal{Q}^n(\omega) e_{k,a}, e_{m,c} \rangle \end{aligned}$$

Hence

$$\begin{aligned} |Vol_{\mathcal{P}}(\{X^{h_{k,a}} : k = 1, 2, \dots, n, a = 1, 2, \dots, d\})| &:= \sqrt{\det \{G^1(X^{h_{k,a}}, X^{h_{m,c}})\}} \\ &= \sqrt{\det \mathcal{Q}^n}. \end{aligned}$$

On the other hand,

$$\begin{aligned} G_{\mathcal{P}}^1(X^{h_{k,a}}, X^{h_{m,c}})(\omega) &= \sum_{i=1}^n g\left(\frac{\nabla X^{h_{k,a}}(\omega, s_{i-1}+)}{ds}, \frac{\nabla X^{h_{m,c}}(\omega, s_{i-1}+)}{ds}\right) \Delta_i s \\ &= \sum_{i=1}^n \langle h'_{k,a}(\omega, s_{i-1}+), h'_{m,c}(\omega, s_{i-1}+) \rangle \Delta_i s \\ &= \sum_{i=1}^n \delta_{k,i} \delta_{m,i} \langle e_a, e_c \rangle \Delta_i s = \delta_{k,m} \delta_{a,c} \Delta_k s, \end{aligned}$$

i.e.  $\{G_{\mathcal{P}}^1(X^{h_{k,a}}, X^{h_{m,c}})\} = \frac{1}{n} \mathcal{I}^n$  and hence

$$\begin{aligned} |Vol_{\mathcal{P}}(\{X^{h_{k,a}} : k = 1, 2, \dots, n, a = 1, 2, \dots, d\})| &:= \sqrt{\det \{G_{\mathcal{P}}^1(X^{h_{k,a}}, X^{h_{m,c}})\}} \\ &= \sqrt{\prod_{i=1}^n (\Delta_i s)^d} = n^{-dn/2}. \end{aligned}$$

Hence it follows that

$$\rho_n \circ \phi = \frac{\sqrt{\det \mathcal{Q}^n}}{\sqrt{\prod_{i=1}^n (\Delta_i s)^d}} = \sqrt{\det(n\mathcal{Q}^n)}.$$

■

Our next goal is to estimate the size of  $\det(n\mathcal{Q}^n)$ . For this we are going to apply Proposition C.2 in the Appendix as follows. For  $\alpha \geq 1$ , to be chosen later, we have, from Equation (C.4) with  $M = n\mathcal{Q}^n$ , that

$$\begin{aligned} \det(n\mathcal{Q}^n) &\leq \alpha^{nd} \exp(\alpha^{-1} \text{Tr}(n\mathcal{Q}^n - \mathcal{I}^n)) = \alpha^{nd} \exp\left(\alpha^{-1} \sum_{m=1}^n \text{tr}(n\mathcal{Q}_{mm}^n - I)\right) \\ &\leq \alpha^{nd} \exp\left(\alpha^{-1} d \sum_{m=1}^n \|n\mathcal{Q}_{mm}^n - I\|\right), \end{aligned} \tag{3.3}$$

where we have used Equation (C.2) of Proposition C.1 in the last inequality. So according to Equation (3.3) we need to estimate  $\|n\mathcal{Q}_{mm}^n - I\|$  for each  $m$ . The first step in the process is to record a formula for  $\mathcal{Q}_{mm}^n$ .

### 3.2 A Formula for $\mathcal{Q}^n$

**Notation 3.5** Given any partition,  $\mathcal{P} := \{0 = s_0 < s_1 < \dots < s_n = 1\}$  of  $[0, 1]$ ,  $j \in \{1, 2, \dots, n\}$ , and  $s \in [0, 1]$ , let

$$[s]_j := [(s - s_{j-1}) \vee 0] \wedge \Delta_j s = \begin{cases} 0 & \text{if } s \leq s_{j-1} \\ s - s_{j-1} & \text{if } s \in J_j \\ \Delta_j s & \text{if } s \geq s_j \end{cases}.$$

**Definition 3.6** For  $i = 1, 2, \dots, n$  and  $0 \leq s \leq \Delta_i s$ , let

$$A_i(\omega, s) := \Omega_{u(s_{i-1}+s)}(\omega'_i, \cdot) \omega'_i \quad (3.4)$$

and  $C_i(\omega, s)$  and  $S_i(\omega, s)$  be the  $\text{End}(\mathbb{R}^d)$ -valued functions determined by

$$\frac{d^2 C_i(\omega, s)}{ds^2} = A_i(\omega, s) C_i(\omega, s) \text{ with } C_i(\omega, 0) = I \text{ and } C'_i(\omega, 0) = 0$$

and

$$\frac{d^2 S_i(\omega, s)}{ds^2} = A_i(\omega, s) S_i(\omega, s) \text{ with } S_i(\omega, 0) = 0 \text{ and } S'_i(\omega, 0) = I.$$

Let  $K > 0$  be a constant such that

$$\begin{aligned} \|A_i(\omega, s) \Delta_i s^2\| &= \|\Omega_{u(s_{i-1}+s)}(\omega'_i, \cdot) \omega'_i\| \Delta_i s^2 \\ &\leq K \|\Delta_i \omega\|^2. \end{aligned}$$

Refer to Section D of Appendix for the existence of such a  $K$ .

Using this notation it follows that  $h(\omega, \cdot)$  is a continuous function which solves Equation (3.1) on  $[0, 1] \setminus \mathcal{P}_n$ , then

$$h(\omega, s) = C_i(\omega, s - s_{i-1}) h(\omega, s_{i-1}) + S_i(\omega, s - s_{i-1}) h'(\omega, s_{i-1}+) \text{ when } s \in J_i.$$

**Notation 3.7** For  $m, l \in \{1, 2, \dots, n\}$  and  $d \times d$  matrices,  $\{M_k\}_{k=1}^n$ , let

$$\prod_{k=l}^m M_k := \begin{cases} I & \text{if } m < l \\ M_m M_{m-1} \dots M_{l+1} M_l & \text{if } m \geq l \end{cases}.$$

With all of this notation, we may write  $h_{i,a}(\omega, s)$  as in the following lemma.

**Lemma 3.8** *Continuing to use the notation introduced above, we have*

$$h_{m,a}(\omega, s) = \left[ \prod_{k=m+1}^n C_k(\omega, [s]_k) \right] S_m(\omega, [s]_m) e_a \quad (3.5)$$

$$= \begin{cases} 0 & \text{if } s \leq s_{m-1} \\ S_m(\omega, s - s_{m-1}) e_a & \text{if } s \in J_m \\ C_j(\omega, s - s_{j-1}) V_{mj}(\omega) e_a & \text{if } s \in J_j \text{ and } j \geq m+1 \end{cases} \quad (3.6)$$

where

$$V_{mj}(\omega) := \left[ \prod_{k=m+1}^{j-1} C_k(\omega, \Delta_k s) \right] S_m(\omega, \Delta_m s). \quad (3.7)$$

Differentiating Equation (3.6),

$$h'_{m,a}(\omega, s) = 1_{J_m}(s) S'_m(\omega, s - s_{m-1}) e_a + \sum_{j=m+1}^n 1_{J_j}(s) C'_j(\omega, s - s_{j-1}) V_{mj} e_a. \quad (3.8)$$

From Equation (3.8), we learn that

$$\begin{aligned} \langle \mathcal{Q}_{mm}^n(\omega) e_a, e_c \rangle &= \int_{J_m} \langle S'_m(\omega, s - s_{m-1}) e_a, S'_m(\omega, s - s_{m-1}) e_c \rangle ds \\ &+ \sum_{j=m+1}^n \int_{J_j} \langle C'_j(\omega, s - s_{j-1}) V_{mj}(\omega) e_a, C'_j(\omega, s - s_{j-1}) V_{mj}(\omega) e_c \rangle ds \\ &= \left\langle \int_0^{\Delta_m s} S'_m(\omega, s)^T S'_m(\omega, s) ds e_a, e_c \right\rangle \\ &+ \sum_{j=m+1}^n \left\langle V_{mj}^T(\omega) \left[ \int_0^{\Delta_j s} C'_j(\omega, s)^T C'_j(\omega, s) ds \right] V_{mj}(\omega) e_a, e_c \right\rangle \end{aligned}$$

and hence we have shown,

$$\begin{aligned} \mathcal{Q}_{mm}^n(\omega) &= \int_0^{\Delta_m s} S'_m(\omega, s)^T S'_m(\omega, s) ds + \sum_{j=m+1}^n V_{mj}^T(\omega) \left[ \int_0^{\Delta_j s} C'_j(\omega, s)^T C'_j(\omega, s) ds \right] V_{mj}(\omega) \\ &= \int_0^{1/n} S'_m(\omega, s)^T S'_m(\omega, s) ds + \sum_{j=m+1}^n V_{mj}^T(\omega) \left[ \int_0^{1/n} C'_j(\omega, s)^T C'_j(\omega, s) ds \right] V_{mj}(\omega). \end{aligned} \quad (3.9)$$

Noting that

$$\|V_{mj}(\omega)\| = \|V_{mj}^T(\omega)\| \leq \|S_m(\omega, \Delta_m s)\|^2 \prod_{k=m+1}^{j-1} \|C_k(\omega, \Delta_k s)\|^2,$$

it follows from Equation (3.9) that

$$\|n\mathcal{Q}_{mm}^n - I\| \leq A_m + B_m \quad (3.10)$$

where

$$A_m(\omega) := n \int_0^{1/n} \left\| S'_m(\omega, s)^T S'_m(\omega, s) - I \right\| ds \quad (3.11)$$

and

$$\begin{aligned} B_m(\omega) &= \sum_{j=m+1}^n \|S_m(\omega, \Delta_m s)\|^2 \prod_{k=m+1}^{j-1} \|C_k(\omega, \Delta_k s)\|^2 \left\| n \int_0^{1/n} C'_j(\omega, s)^T C'_j(\omega, s) ds \right\| \\ &\leq \sum_{j=m+1}^n \|S_m(\omega, \Delta_m s)\|^2 \prod_{k=m+1}^{j-1} \|C_k(\omega, \Delta_k s)\|^2 n \int_0^{1/n} \|C'_j(\omega, s)\|^2 ds. \end{aligned} \quad (3.12)$$

Thus we are now left to estimate the quantities comprising  $A_m$  and  $B_m$ .

### 3.3 Estimates for Solutions to Jacobi's Equation

**Remark 3.9** *In what follows we will make use of the following elementary estimates without further comment.*

$$1. \cosh x = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!} \leq \sum_{n=0}^{\infty} \frac{x^{2n}}{2^n n!} = e^{x^2/2}.$$

$$2. \cosh x = \cosh |x| = \frac{e^{|x|} + e^{-|x|}}{2} \leq e^{|x|} \text{ so that}$$

$$\cosh x \leq \min \left( e^{x^2/2}, e^{|x|} \right),$$

and

$$3. \text{ for } x \geq 0,$$

$$\sinh x = \int_0^x \cosh t \, dt \leq \int_0^x \cosh x \, dt = x \cosh x.$$

*This estimate is also easily understood using the power series expansions for  $\sinh$  and  $\cosh$ .*

**Lemma 3.10 (Global Estimate)** *Let  $A(s)$  be a  $d \times d$  matrix for all  $s \geq 0$ ,  $\kappa := \sup_{s \geq 0} \|A(s)\| < \infty$ , and let  $Z(s)$  be either a  $\mathbb{R}^d$  or  $d \times d$  matrix valued solution to the second order differential equation*

$$Z''(s) = A(s)Z(s).$$

*Then*

$$\|Z(s) - Z(0)\| \leq \|Z(0)\| (\cosh \sqrt{\kappa} s - 1) + \|Z'(0)\| \frac{\sinh \sqrt{\kappa} s}{\sqrt{\kappa}}. \quad (3.13)$$



**Proof.** By Taylor's theorem with integral remainder,

$$\begin{aligned} Z(s) &= Z(0) + sZ'(0) + \int_0^s Z''(u)(s-u) du \\ &= Z(0) + sZ'(0) + \int_0^s A(u)Z(u)(s-u) du \end{aligned} \quad (3.14)$$

and therefore

$$\begin{aligned} \|Z(s) - Z(0)\| &\leq s \|Z'(0)\| + \kappa \int_0^s \|Z(u)\| (s-u) du \\ &\leq s \|Z'(0)\| + \kappa \int_0^s \|Z(u) - Z(0)\| (s-u) du + \frac{1}{2}s^2\kappa \|Z(0)\| \\ &:= f(s). \end{aligned} \quad (3.15)$$

Note that  $f(0) = 0$ ,

$$f'(s) = \|Z'(0)\| + \kappa \int_0^s \|Z(u) - Z(0)\| (s-u) du + s\kappa \|Z(0)\|,$$

$f'(0) = \|Z'(0)\|$ , and

$$f''(s) = \kappa \|Z(s) - Z(0)\| + \kappa \|Z(0)\| \leq \kappa f(s) + \kappa \|Z(0)\|.$$

That is,

$$f''(s) = \kappa f(s) + \eta(s), \quad f(0) = 0, \quad \text{and} \quad f'(0) = \|Z'(0)\|, \quad (3.16)$$

where  $\eta(s) := f''(s) - \kappa f(s) \leq \kappa \|Z(0)\|$ . Equation (3.16) may be solved by variation of parameters to find

$$\begin{aligned} f(s) &= \|Z'(0)\| \frac{\sinh \sqrt{\kappa}s}{\sqrt{\kappa}} + \int_0^s \frac{\sinh \sqrt{\kappa}(s-r)}{\sqrt{\kappa}} \eta(r) dr \\ &\leq \|Z'(0)\| \frac{\sinh \sqrt{\kappa}s}{\sqrt{\kappa}} + \|Z(0)\| \int_0^s \sqrt{\kappa} \sinh \sqrt{\kappa}(s-r) dr \\ &= \|Z'(0)\| \frac{\sinh \sqrt{\kappa}s}{\sqrt{\kappa}} + \|Z(0)\| (\cosh \sqrt{\kappa}s - 1). \end{aligned}$$

Combining this with Equation (3.15) proves Equation (3.13). ■

**Theorem 3.11** Suppose that  $A(s)$  above satisfies,  $0 \leq -A(s) \leq \kappa I$  for all  $s$  or equivalently that  $-\kappa I \leq A(s) \leq 0$  for all  $s$ . Let

$$\psi(s) := \min \left( 1 + \frac{\cosh(s)}{16} s^4, \cosh(s) \right)$$

whose graph is shown below and  $C(s)$  and  $S(s)$  be the matrix functions defined by

$$\begin{aligned} C''(s) &= A(s)C(s), \text{ with } C(0) = I, \quad C'(0) = 0, \\ S''(s) &= A(s)S(s), \text{ with } S(0) = 0, \quad S'(0) = I. \end{aligned}$$




Figure 1: Graph of  $\psi$ .

Then

1.  $\|C(s)\| \leq \psi(\sqrt{\kappa}s)$ ,
2.  $\|S(s)\| \leq s\psi(\sqrt{\kappa}s)$ ,
3.  $\|C'(s)\| \leq \kappa s\psi(\sqrt{\kappa}s)$ ,
4.  $\|S'(s)\| \leq 1 + \frac{1}{2}\kappa s^2\psi(\sqrt{\kappa}s)$ , and
5.  $\left\|S'(s)^T S'(s) - I\right\| \leq \psi(\sqrt{\kappa}s)\kappa s^2 + \frac{1}{3}\psi^2(\sqrt{\kappa}s)\kappa^2 s^4$ .

Moreover, if we only assume that  $\|A(s)\| \leq \kappa$ , all of the above estimates still hold provided that  $\psi$  is replaced by  $\cosh$ .

**Proof.** If  $Z$  solves,  $Z''(s) = A(s)Z(s)$ , then iterating Equation (3.14) shows

$$\begin{aligned}
 Z(s) &= Z(0) + sZ'(0) + \int_0^s A(u) \left[ Z(0) + uZ'(0) + \int_0^u A(r)Z(r)(u-r) dr \right] (s-u) du \\
 &= \left( I + \int_0^s (s-u)A(u)du \right) Z(0) + \left( sI + \int_0^s u(s-u)A(u) du \right) Z'(0) \\
 &\quad + \int_{0 \leq r \leq u \leq s} (u-r)(s-u)A(u)A(r)Z(r)drdu.
 \end{aligned}$$

In particular this shows

$$C(s) = \left( I - \int_0^s (s-u)[-A(u)] du \right) + \int_{0 \leq r \leq u \leq s} (u-r)(s-u)A(u)A(r)C(r)drdu \quad (3.17)$$

and

$$S(s) = \left( sI - \int_0^s u(s-u) [-A(u)] du \right) + \int_{0 \leq r \leq u \leq s} (u-r)(s-u) A(u) A(r) S(r) dr du. \quad (3.18)$$

From Equation (3.17),

$$\begin{aligned} \|C(s)\| &\leq \left\| I + \int_0^s (s-u) A(u) du \right\| + \int_{0 \leq r \leq u \leq s} (u-r)(s-u) \|A(u) A(r) C(r)\| dr du \\ &\leq \left\| I + \int_0^s (s-u) A(u) du \right\| + \frac{\cosh(\sqrt{\kappa}s)}{24} \kappa^2 s^4. \end{aligned}$$

Moreover,

$$\left(1 - \kappa \frac{s^2}{2}\right) I = I - \kappa \left[ \int_0^s (s-u) du \right] I \leq I + \int_0^s (s-u) A(u) du \leq I$$

from which it follows that

$$\left\| I + \int_0^s (s-u) A(u) du \right\| \leq \max \left( 1, \kappa \frac{s^2}{2} - 1 \right) = 1_{s^2 \leq 4/\kappa} + \left( \kappa \frac{s^2}{2} - 1 \right) 1_{s^2 \geq 4/\kappa}$$

and hence we have

$$\|C(s)\| \leq 1_{s^2 \leq 4/\kappa} + \left( \kappa \frac{s^2}{2} - 1 \right) 1_{s^2 \geq 4/\kappa} + \frac{\cosh(\sqrt{\kappa}s)}{24} \kappa^2 s^4 \leq 1 + \frac{\cosh(\sqrt{\kappa}s)}{16} \kappa^2 s^4.$$

This is because  $f(s) = s^4 \frac{\cosh s}{48} - \left( \frac{s^2}{2} - 1 \right) = 1 + s^2 \left[ s^2 \frac{\cosh s}{48} - \frac{1}{2} \right]$  is an increasing function and for  $s \geq 2$ ,  $f(s) \geq f(2) > 0.25$ . Recall that we also know that  $\|C(s)\| \leq \cosh(\sqrt{\kappa}s)$  and therefore we have

$$\begin{aligned} \|C(s)\| &\leq \min \left( 1_{s^2 \leq 4/\kappa} + \left( \kappa \frac{s^2}{2} - 1 \right) 1_{s^2 \geq 4/\kappa} + \frac{\cosh(\sqrt{\kappa}s)}{24} \kappa^2 s^4, \cosh(\sqrt{\kappa}s) \right) \\ &\leq \min \left( 1 + \frac{\cosh(\sqrt{\kappa}s)}{16} \kappa^2 s^4, \cosh(\sqrt{\kappa}s) \right). \end{aligned}$$

It will be convenient to define

$$\psi(s) := \min \left( 1 + \frac{\cosh(s)}{16} s^4, \cosh(s) \right)$$

and hence

$$\|C(s)\| \leq \psi(\sqrt{\kappa}s).$$

Similarly, from Equation (3.18)

$$\begin{aligned} \|S(s)\| &\leq \left\| sI + \int_0^s u(s-u) A(u) du \right\| + \int_{0 \leq r \leq u \leq s} (u-r)(s-u) \|A(u) A(r) S(r)\| dr du \\ &\leq s \left\| I + \frac{1}{s} \int_0^s u(s-u) A(u) du \right\| + \frac{\kappa^2}{24} s^4 \frac{\sinh(\sqrt{\kappa}s)}{\sqrt{\kappa}}. \end{aligned}$$

In this case,

$$\left(1 - \frac{\kappa}{6} s^2\right) I = \left(1 - \kappa \frac{1}{s} \int_0^s u(s-u) du\right) I \leq I + \frac{1}{s} \int_0^s u(s-u) A(u) du \leq I.$$

Combining this with the previous equation shows

$$\begin{aligned}\|S(s)\| &\leq s \left[ 1_{s^2 \leq 12/\kappa} + \left( \kappa \frac{s^2}{6} - 1 \right) 1_{s^2 \geq 12/\kappa} \right] + \frac{\kappa^2}{24} s^4 \frac{\sinh(\sqrt{\kappa}s)}{\sqrt{\kappa}} \\ &\leq s \left[ 1_{s^2 \leq 12/\kappa} + \left( \kappa \frac{s^2}{6} - 1 \right) 1_{s^2 \geq 12/\kappa} + \frac{\cosh(\sqrt{\kappa}s)}{24} \kappa^2 s^4 \right] \\ &\leq s \left[ 1 + \frac{\cosh(\sqrt{\kappa}s)}{16} \kappa^2 s^4 \right].\end{aligned}$$

Since we also have

$$\|S(s)\| \leq \frac{\sinh(\sqrt{\kappa}s)}{\sqrt{\kappa}} \leq s \cosh(\sqrt{\kappa}s),$$

we may conclude that

$$\|S(s)\| \leq s \min \left( 1 + \frac{\cosh(\sqrt{\kappa}s)}{16} \kappa^2 s^4, \cosh(\sqrt{\kappa}s) \right) = s\psi(\sqrt{\kappa}s).$$

Furthermore,

$$\|C'(s)\| = \left\| \int_0^s A(r) C(r) dr \right\| \leq \int_0^s \|A(r)\| \|C(r)\| dr \leq \kappa s \psi(\sqrt{\kappa}s)$$

and

$$\begin{aligned}\|S'(s)\| &= \left\| I + \int_0^s A(r) S(r) dr \right\| \leq 1 + \int_0^s \|A(r)\| \|S(r)\| dr \\ &\leq 1 + \kappa \int_0^s r \psi(\sqrt{\kappa}r) dr \leq 1 + \frac{1}{2} \kappa s^2 \psi(\sqrt{\kappa}s).\end{aligned}$$

Finally

$$\frac{d}{ds} [S'(s)^T S'(s)] = S(s)^T A(s) S'(s) + S'(s)^T A(s) S(s)$$

and therefore,

$$\begin{aligned}\left\| \frac{d}{ds} [S'(s)^T S'(s)] \right\| &\leq 2 \|S(s)\| \|A(s)\| \|S'(s)\| \\ &\leq 2 \|S(s)\| \|A(s)\| \|S'(s)\| \\ &\leq 2 \kappa s \psi(\sqrt{\kappa}s) \left( 1 + \frac{1}{2} \kappa s^2 \psi(\sqrt{\kappa}s) \right).\end{aligned}$$

Integrating this equation then implies,

$$\begin{aligned}\|S'(s)^T S'(s) - I\| &\leq \int_0^s 2 \kappa r \psi(\sqrt{\kappa}r) \left( 1 + \frac{1}{2} \kappa r^2 \psi(\sqrt{\kappa}r) \right) dr \\ &\leq \int_0^s 2 \kappa \psi(\sqrt{\kappa}s) \left( r + \frac{1}{2} \kappa r^3 \psi(\sqrt{\kappa}s) \right) dr \\ &= \kappa \psi(\sqrt{\kappa}s) \left( s^2 + \frac{1}{3} \kappa s^4 \psi(\sqrt{\kappa}s) \right) \\ &= \psi(\sqrt{\kappa}s) \kappa s^2 + \frac{1}{3} \psi^2(\sqrt{\kappa}s) \kappa^2 s^4.\end{aligned}$$

■

**Proposition 3.12** *If  $\|A(s)\| \leq \kappa$ , then the following estimates hold:*

$$\left\| S'(s) - \left( I + \int_0^s r A(r) dr \right) \right\| \leq s^4 \kappa^2 \cosh(\sqrt{\kappa} s) \quad (3.19)$$

$$\left\| \frac{S(s)}{s} - \left( I + \frac{1}{s} \int_0^s (s-r) r A(r) dr \right) \right\| \leq s^4 \kappa^2 \cosh(\sqrt{\kappa} s) \quad (3.20)$$

$$\left\| C(s) - \left( I + \int_0^s (s-u) A(u) du \right) \right\| \leq s^4 \kappa^2 \cosh(\sqrt{\kappa} s). \quad (3.21)$$

**Proof.** Now

$$\begin{aligned} S'(s) &= I + \int_0^s A(r) S(r) dr = I + \int_0^s A(r) \left[ \int_0^r S'(u) du \right] dr \\ &= I + \int_0^s A(r) \left[ \int_0^r \left[ I + \int_0^u A(v) S(v) dv \right] du \right] dr \\ &= I + \int_0^s r A(r) dr + \int_{0 \leq v \leq u \leq r \leq s} A(r) A(v) S(v) dv du dr \\ &= I + \int_0^s r A(r) dr + \int_{0 \leq v \leq r \leq s} (r-v) A(r) A(v) S(v) dv dr. \end{aligned}$$

Thus

$$\begin{aligned} \left\| S'(s) - \left( I + \int_0^s r A(r) dr \right) \right\| &\leq \int_{0 \leq v \leq r \leq s} (r-v) \kappa^2 v \cosh(\sqrt{\kappa} v) dv dr \\ &\leq s^4 \kappa^2 \cosh(\sqrt{\kappa} s), \end{aligned}$$

Integrating this estimate implies

$$\left\| S(s) - Is - \int_0^s (s-r) A(r) dr \right\| \leq s \cdot s^4 \kappa^2 \cosh(\sqrt{\kappa} s)$$

which is equivalent to Equation (3.20). Similarly from Equation (3.17) we have

$$\begin{aligned} \left\| C(s) - \left( I + \int_0^s (s-u) A(u) du \right) \right\| &= \left\| \int_{0 \leq r \leq u \leq s} (u-r)(s-u) A(u) A(r) C(r) dr du \right\| \\ &\leq s^4 \kappa^2 \cosh(\sqrt{\kappa} s). \end{aligned}$$

■

**Definition 3.13** *Let*

$$h(t) = \frac{1}{t} \ln \psi(\sqrt{t}) = t^{-1} \ln \left( \min \left( 1 + \frac{\cosh(\sqrt{t})}{16} t^2, \cosh(\sqrt{t}) \right) \right)$$

*whose graph is given in Figure 2, i.e.*

$$e^{s^2 h(s^2)} = \psi(s).$$




Figure 2: Here is the graph of  $h(t)$ .

Let  $\varphi$  be a function, which we will specify shortly, such that  $\varphi \geq \psi$ . Further let  $g(t) := \frac{1}{t} \ln \varphi(\sqrt{t})$  so that  $\varphi(s) = e^{s^2 g(s^2)}$  and define

$$u(t) := \frac{\psi^2(\sqrt{t})}{\varphi^2(\sqrt{t})} = \frac{e^{2th(t)}}{e^{2tg(t)}} = e^{-2t(g(t)-h(t))}.$$

We will specify  $\varphi(t)$  by requiring  $g(t)$  to be a smooth function such that  $g(t) = h(t)$  for  $t$  near zero which then rises rapidly to a height of .6 as  $t$  increases. For later purposes, let us observe that with this definition  $tu(t)$  is bounded by .63 as the graph below indicates.



Figure 3: Graph of  $tu(t)$  and the line  $y = .63$ . We see that  $tu(t) \leq .63$ .

### 3.4 Proof of Uniform Integrability

**Proposition 3.14** *Suppose  $N$  is a  $d$  - dimensional standard Gaussian normal random variable and  $G(x, \|N\|)$  is a  $C^1$  - function in  $x \in (-\varepsilon, \varepsilon)$  such that  $G(0, \|N\|) \equiv 0$ ,*

$$\alpha := \sup \{G(x, t) : x \in (-\varepsilon, \varepsilon) \text{ and } t \geq 0\} < \frac{1}{2}$$

*and there exist constants  $C < \infty$  and  $\beta < \infty$  such that*

$$G_x(x, t) = \frac{\partial G}{\partial x}(x, t) \leq C(1+t)^\beta \text{ for } x \in (-\varepsilon, \varepsilon) \text{ and } t \geq 0.$$

*Then*

$$\lim_{n \rightarrow \infty} \left( \mathbb{E} \left[ e^{\|N\|^2 G(\frac{1}{n}, \|N\|)} \right] \right)^n = e^{\mathbb{E}[\|N\|^2 G_x(0, \|N\|)]}.$$

**Proof.** Let

$$f(x) := \mathbb{E} \left[ e^{\|N\|^2 G(x, \|N\|)} \right].$$

Since

$$e^{\|N\|^2 G(x, \|N\|)} \leq e^{\alpha \|N\|^2}$$

and

$$G_x(x, \|N\|) \|N\|^2 e^{\|N\|^2 G(x, \|N\|)} \leq C(1 + \|N\|)^\beta \|N\|^2 e^{\alpha \|N\|^2}$$

with the right hand members of these inequalities being integrable functions, it follows that  $f(x)$  is a  $C^1$  - function for  $x$  near 0 with

$$f'(x) = \mathbb{E} \left[ G_x(x, \|N\|) \|N\|^2 e^{\|N\|^2 G(x, \|N\|)} \right].$$

In particular we have  $f(0) = \mathbb{E} \left[ e^{\|N\|^2 G(0, \|N\|)} \right] = 1$  and

$$f'(0) = \mathbb{E} \left[ G_x(0, \|N\|) \|N\|^2 \right]$$

Therefore,

$$\begin{aligned} \lim_{n \rightarrow \infty} \left( \mathbb{E} \left[ e^{\|N\|^2 G(\frac{1}{n}, \|N\|)} \right] \right)^n &= \lim_{n \rightarrow \infty} (f(n^{-1}))^n = \lim_{n \rightarrow \infty} e^{n \ln f(n^{-1})} = \lim_{x \rightarrow 0} e^{\frac{1}{x} \ln f(x)} \\ &= \lim_{x \rightarrow 0} e^{f'(x)/f(x)} = e^{f'(0)} = e^{\mathbb{E}[G_x(0, \|N\|) \|N\|^2]}. \end{aligned}$$

■

**Theorem 3.15** *Suppose that  $(M, g)$  is a Riemannian manifold with non-negative sectional curvatures which are bounded above by  $K = \frac{1}{2d}$ . Then for all  $p$  sufficiently close to 1,*

$$\sup_n \mathbb{E} \left[ \det^{p/2}(n\mathcal{Q}^n \circ b_n) \right] < \infty.$$

**Proof.** For  $\omega \in H_{\mathcal{P}}(\mathbb{R}^d)$ , from Theorem 3.11,

$$\begin{aligned} A_m(\omega) &= n \int_0^{1/n} \left\| S'_m(\omega, s)^T S'_m(\omega, s) - I \right\| ds \\ &\leq \psi \left( \sqrt{K} \|\Delta_m \omega\| \right) K \|\Delta_m \omega\|^2 + \frac{1}{3} \psi^2 \left( \sqrt{K} \|\Delta_m \omega\| \right) K^2 \|\Delta_m \omega\|^4, \end{aligned}$$

and if we write  $\tau_j(\omega) = \sqrt{K} \|\Delta_j \omega\|$ ,

$$\begin{aligned} B_m(\omega) &= \sum_{j=m+1}^n \|S_m(\omega, \Delta_m s)\|^2 \prod_{k=m+1}^{j-1} \|C_k(\omega, \Delta_k s)\|^2 n \int_0^{1/n} \|C'_j(\omega, s)\|^2 ds \\ &\leq \sum_{j=m+1}^n (\Delta_m s)^2 \psi^2(\tau_m(\omega)) \left[ \prod_{k=m+1}^{j-1} \psi^2(\tau_k(\omega)) \right] K^2 (\Delta_j s)^2 \|\omega'_j\|^4 \psi^2(\tau_j(\omega)) \\ &= K^2 \sum_{j=m+1}^n \left[ \prod_{k=m}^j \psi^2(\tau_k(\omega)) \right] \|\Delta_j \omega\|^4 \\ &= K^2 \sum_{j=m+1}^n \left[ \prod_{k=m}^j \psi^2 \left( \sqrt{K} \|\Delta_k \omega\| \right) \right] \|\Delta_j \omega\|^4. \end{aligned}$$

Hence if we let

$$\alpha(\omega) := \prod_{k=1}^n \varphi^2 \left( \sqrt{K} \|\Delta_k \omega\| \right) = \prod_{k=1}^n e^{2K \|\Delta_k \omega\|^2 g(K \|\Delta_k \omega\|^2)},$$

then

$$\begin{aligned} A_m(\omega) &\leq \alpha(\omega) K \|\Delta_m \omega\|^2 + \frac{1}{3} \alpha(\omega) u \left( K \|\Delta_m \omega\|^2 \right) K^2 \|\Delta_m \omega\|^4 \\ &\leq \alpha(\omega) \left[ K \|\Delta_m \omega\|^2 + \frac{1}{3} u \left( K \|\Delta_m \omega\|^2 \right) K^2 \|\Delta_m \omega\|^4 \right] \end{aligned}$$

and

$$\begin{aligned} B_m(\omega) &\leq \alpha(\omega) K^2 \sum_{j=m+1}^n \left[ \prod_{k=m}^j u \left( \sqrt{K} \|\Delta_k \omega\| \right) \right] \|\Delta_j \omega\|^4 \\ &\leq \alpha(\omega) K^2 \sum_{j=1}^n u \left( \sqrt{K} \|\Delta_j \omega\| \right) \|\Delta_j \omega\|^4. \end{aligned}$$

For  $\omega \in (W(\mathbb{R}^d), \mu)$ ,

$$\Delta_i b_n(\omega) = \Delta_i b(\omega) = b(s_i)(\omega) - b(s_{i-1})(\omega) = \omega(s_i) - \omega(s_{i-1}) = \Delta_i \omega.$$



Therefore, on  $W(\mathbb{R}^d)$ ,

$$\begin{aligned}
& \sum_{m=1}^n (A_m + B_m) \circ b_n \\
& \leq \alpha \circ b_n \cdot \left[ \sum_{m=1}^n \left[ K \|\Delta_m b\|^2 + \frac{1}{3} u \left( K \|\Delta_m b\|^2 \right) K^2 \|\Delta_m b\|^4 \right] \right. \\
& \quad \left. + K^2 n \sum_{m=1}^n u \left( K \|\Delta_m b\|^2 \right) \|\Delta_m b\|^4 \right] \\
& = \alpha \circ b_n \cdot \sum_{m=1}^n \left[ K \|\Delta_m b\|^2 + \left( \frac{1}{3} + n \right) K^2 u \left( \sqrt{K} \|\Delta_m b\| \right) \|\Delta_m b\|^4 \right].
\end{aligned}$$

Now let  $x = n^{-1}$  and  $N_m := \sqrt{n} \Delta_m b$ , so that  $\{N_m\}_{m=1}^n$  is a collection of  $\mathbb{R}^d$ -valued independent standard normal random variables. With this notation we have

$$\begin{aligned}
& \sum_{m=1}^n (A_m + B_m) \circ b_n \\
& \leq \alpha \circ b_n \cdot \sum_{m=1}^n \left[ Kx \|N_m\|^2 + \left( \frac{1}{3} + n \right) K^2 u \left( Kx \|N_m\|^2 \right) x^2 \|N_m\|^4 \right] \\
& = \alpha \circ b_n \cdot \sum_{m=1}^n \left[ Kx \|N_m\|^2 + K \left( \frac{1}{3}x + 1 \right) u \left( Kx \|N_m\|^2 \right) Kx \|N_m\|^4 \right] \\
& = \alpha \circ b_n \cdot \sum_{m=1}^n \left[ Kx + K \left( \frac{1}{3}x + 1 \right) u \left( Kx \|N_m\|^2 \right) Kx \|N_m\|^2 \right] \|N_m\|^2
\end{aligned}$$

and therefore, using Equation (3.3),

$$\begin{aligned}
& \det(n\mathcal{Q}^n \circ b_n) = \rho_n \circ \phi \circ b_n \\
& \leq \left( \alpha^{nd} \exp \left( \alpha^{-1} d \sum_{m=1}^n \|(n\mathcal{Q}_{mm}^n - I)\| \right) \right) \circ b_n \\
& \leq \left( \alpha^{nd} \exp \left( \alpha^{-1} d \sum_{m=1}^n (A_m + B_m) \right) \right) \circ b_n \\
& \leq (\alpha \circ b_n)^{nd} \exp \left( d \sum_{m=1}^n \left[ Kx + K \left( \frac{1}{3}x + 1 \right) u \left( Kx \|N_m\|^2 \right) Kx \|N_m\|^2 \right] \|N_m\|^2 \right)
\end{aligned}$$

where

$$\begin{aligned}
(\alpha \circ b_n)^{nd} &= \prod_{m=1}^n \varphi^{2nd} \left( \sqrt{K} \|\Delta_m b\| \right) = \prod_{m=1}^n \varphi^{2nd} \left( \sqrt{Kx} \|N_m\| \right) \\
&= \prod_{m=1}^n e^{2ndK \|\Delta_m b\|^2 g(K \|\Delta_m b\|^2)} = \prod_{m=1}^n e^{2dK \|N_m\|^2 g(Kx \|N_m\|^2)}.
\end{aligned}$$

Let  $\theta(x, \|N\|^2) = x + (\frac{1}{3}x + 1)u(Kx\|N\|^2)Kx\|N\|^2$ . Then,

$$\begin{aligned}\det(n\mathcal{Q}^n \circ b_n) &\leq \prod_{m=1}^n e^{2dK\|N_m\|^2 g(Kx\|N_m\|^2)} \exp\left(dK\theta(x, \|N_m\|^2)\|N_m\|^2\right) \\ &= \prod_{m=1}^n \exp\left(dK\left[2g(Kx\|N_m\|^2) + \theta(x, \|N_m\|^2)\right]\|N_m\|^2\right).\end{aligned}$$

Hence

$$\begin{aligned}\mathbb{E}\left[\det^{p/2}(n\mathcal{Q}^n \circ b_n)\right] &= \left[\mathbb{E}\exp\left(dpK\left[g(Kx\|N\|^2) + \frac{1}{2}\theta(x, \|N\|^2)\right]\|N\|^2\right)\right]^n \\ &= \left[\mathbb{E}\exp\left(dpK \cdot G(x, \|N\|^2)\|N\|^2\right)\right]^{1/x}\end{aligned}$$

where

$$G(x, \|N\|^2) = g(Kx\|N\|^2) + \frac{x}{2} + \left(\frac{1}{6}x + \frac{1}{2}\right)u(Kx\|N\|^2)Kx\|N\|^2.$$

By our choice of  $g$  and hence  $u$ , we know

$$g(Kx\|N\|^2) + \frac{1}{2}u(Kx\|N\|^2)Kx\|N\|^2 \leq .6 + \frac{1}{2}.63 = .915 < 1.$$

Therefore, for small  $x$ ,  $G(x, \|N\|^2) \leq .92 < 1$  for small  $x$ . Hence, if  $p$  is sufficiently close to 1, we will have

$$dpK \cdot G(x, \|N\|^2) = \frac{1}{2}pG(x, \|N\|^2) \leq \frac{1}{2}p \cdot 0.92 < \frac{1}{2}.$$

Therefore we may apply Proposition 3.14 to conclude that

$$\begin{aligned}\limsup_{n \rightarrow \infty} \mathbb{E}\left[\det^{p/2}(n\mathcal{Q}^n \circ b_n)\right] &\leq \lim_{x \rightarrow 0} \left[\mathbb{E}\exp\left(dpK \cdot G(x, \|N\|^2)\|N\|^2\right)\right]^{1/x} \\ &= \exp\left(\mathbb{E}\left[dpK \cdot G_x(0, \|N\|^2)\|N\|^2\right]\right) \\ &= \exp\left(\mathbb{E}\left[dpK\left(\frac{2K}{16}\|N\|^2 + \frac{1}{2} + \frac{1}{2}K\|N\|^2\right)\|N\|^2\right]\right) < \infty.\end{aligned}$$

■

## 4 Second Formula for $\rho_n$

**Definition 4.1** For any  $\epsilon > 0$  and any partition  $\mathcal{P}$  of  $[0, 1]$ , let

$$\begin{aligned}H_{\mathcal{P}}^{\epsilon}(\mathbb{R}^d) &= \left\{\omega \in H_{\mathcal{P}}(\mathbb{R}^d) \mid \int_{s_{i-1}}^{s_i} \|\omega'(s)\| ds < \epsilon \text{ for } i = 1, \dots, n\right\} \\ &= \{\omega \in H_{\mathcal{P}}(\mathbb{R}^d) \mid \|\Delta_i \omega\| < \epsilon \forall i\},\end{aligned}$$

where  $\Delta_i \omega = \omega(s_i) - \omega(s_{i-1})$ . The second equality holds since  $\omega'_i(s)$  is a constant in  $J_i$  for each  $i$ .

**Remark 4.2** We will now consider all the  $\omega \in H_{\mathcal{P}}^\epsilon(\mathbb{R}^d)$ , with  $\epsilon$  sufficiently small, specified in the next lemma.

**Lemma 4.3** There exists an  $\epsilon$  with  $\bigvee_{i=1,\dots,n} \|\Delta_i \omega\| < \epsilon$  such that for  $i = 1, 2, \dots, n$ ,  $S_i(\omega, s)$  is invertible for  $0 < s \leq \Delta_i s$ .

**Proof.** From Equation (3.13) with  $\kappa = \frac{K}{(\Delta_i s)^2} \|\Delta_i \omega\|^2$ , we see that

$$\|S_i(\omega, s)\| \leq s \left( \frac{\sinh(\sqrt{K} \|\Delta_i \omega\|)}{\sqrt{K} \|\Delta_i \omega\|} \right) \leq s \cosh(\sqrt{K} \|\Delta_i \omega\|), \quad (4.1)$$

where we have used the inequality  $\frac{\sinh x}{x} \leq \cosh x$ . By Taylor's Theorem with integral remainder,

$$S_i(\omega, s) = sI + \int_0^s (s-u) S_i^{(2)}(\omega, u) du = sI + \int_0^s (s-u) A_i(\omega, u) S_i(\omega, u) du.$$

Now using Theorem 3.11,

$$\begin{aligned} \left\| \int_0^s (s-u) A_i(\omega, u) S_i(\omega, u) du \right\| &\leq \int_0^s (s-u) \frac{K \|\Delta_i \omega\|^2}{(\Delta_i s)^2} u \cosh(\sqrt{K} \|\Delta_i \omega\|) du \\ &= \left( \int_0^s (s-u) u du \right) \frac{K \|\Delta_i \omega\|^2}{(\Delta_i s)^2} \cosh(\sqrt{K} \|\Delta_i \omega\|) \\ &= \frac{s^3}{6} \frac{K}{(\Delta_i s)^2} \|\Delta_i \omega\|^2 \cosh(\sqrt{K} \|\Delta_i \omega\|) \\ &\leq sK \|\Delta_i \omega\|^2 \cosh(\sqrt{K} \|\Delta_i \omega\|). \end{aligned}$$

Hence, if we choose an  $\epsilon$  such that for  $0 < x \leq \epsilon$ ,

$$Kx^2 \cosh(\sqrt{K}x) < 1,$$

then

$$S_i(\omega, s) = s \left( I + \frac{1}{s} \int_0^s (s-u) A_i(\omega, u) S_i(\omega, u) du \right)$$

is invertible for  $0 < s \leq \Delta_i s$ . ■

In order to compute  $\lim_{n \rightarrow 0} \rho_n$ , we will first derive another formula for  $\rho_n$ . Define a set of tangent vectors  $\{f_{i,a}(\omega, s)\}_{\substack{i=1,2,\dots,n \\ a=1,2,\dots,d}}$  on  $T_\omega H_{\mathcal{P}}^\epsilon(\mathbb{R}^d)$  such that  $f_{i,a}(\omega, s)$  is the solution to Equation (3.1) with the given initial conditions

$$f_{i,a}(\omega, 0) = 0,$$

for  $j = 1, \dots, n$ ,

$$f'_{i,a}(\omega, s_{j-1}) = \begin{cases} e_a, & j = i \\ -F_i(\omega) e_a, & j = i + 1 \\ 0, & \text{otherwise.} \end{cases}$$

where

$$F_i(\omega) := (S_{i+1}(\omega, \Delta_{i+1}s))^{-1} C_{i+1}(\omega, \Delta_{i+1}s) S_i(\omega, \Delta_i s), \quad (4.2)$$

where  $S_i$  and  $C_i$  are as in Definition 3.6. By Lemma 4.3, we can choose an  $\epsilon$  such that  $F_i(\omega)$  is defined on  $H_{\mathcal{P}}^\epsilon(\mathbb{R}^d)$ . Therefore,

$$f_{i,a}(\omega, s) = \begin{cases} S_i(\omega, s - s_{i-1}) e_a, & s \in J_i \\ C_{i+1}(\omega, s - s_i) S_i(\omega, \Delta_i s) e_a - S_{i+1}(\omega, s - s_i) F_i(\omega) e_a, & s \in J_{i+1} \\ 0, & \text{otherwise,} \end{cases}$$

and hence

$$f'_{i,a}(\omega, s) = \begin{cases} F_{ii}(\omega, s) e_a, & s \in J_i \\ F_{i+1,i}(\omega, s) e_a, & s \in J_{i+1} \\ 0, & \text{otherwise,} \end{cases}$$

where for  $i = 1, \dots, n$ ,

$$\begin{aligned} F_{ii}(\omega, s) &= S'_i(\omega, s - s_{i-1}), \\ F_{i+1,i}(\omega, s) &= C'_{i+1}(\omega, s - s_i) S_i(\omega, \Delta_i s) - S'_{i+1}(\omega, s - s_i) F_i(\omega). \end{aligned}$$

**Remark 4.4** For all  $\sigma \in H_{\mathcal{P}}^\epsilon(M)$ , the vectors,  $\{X^{f_{i,a}}(\sigma)\}_{i=1, \dots, n, a=1, \dots, d}$  form a basis for  $T_\sigma H_{\mathcal{P}}^\epsilon(M)$ , where  $\sigma = \phi(\omega)$ .

At this point, we will now assume  $\mathcal{P}_n = \{s_i = \frac{i}{n}\}_{i=0}^n$  and  $\Delta = 1/n$  throughout this section. We may now write

$$f'_{i,a}(\omega, s) = [1_{J_i}(s) S'_i(\omega, s - s_{i-1}) + 1_{J_{i+1}}(s) V'_{i+1}(\omega, s - s_i)] e_a \quad (4.3)$$

where  $V_1 \equiv 0 \equiv V_{n+1}$  and for  $2 \leq i \leq n$ ,

$$V_i(\omega, s) := C_i(\omega, s) S_{i-1}(\omega, \Delta) - S_i(\omega, s) F_{i-1}(\omega)$$

and

$$F_i(\omega) := S_{i+1}(\omega, \Delta)^{-1} C_{i+1}(\omega, \Delta) S_i(\omega, \Delta). \quad (4.4)$$

Observe that

$$V_i(\omega, \Delta) = C_i(\omega, \Delta) S_{i-1}(\omega, \Delta) - S_i(\omega, \Delta) S_i(\omega, \Delta)^{-1} C_i(\omega, \Delta) S_{i-1}(\omega, \Delta) = 0 \quad (4.5)$$

and that

$$V_i(\omega, 0) = S_{i-1}(\omega, \Delta). \quad (4.6)$$

**Lemma 4.5** *Continuing the notation above, we have*

$$\begin{aligned}
& \int_0^1 \langle f'_{i,a}(\omega, s), f'_{j,c}(\omega, s) \rangle ds \\
&= \begin{cases} \int_0^\Delta \langle S'_i(\omega, s) e_a, V'_i(\omega, s) e_c \rangle ds & \text{if } j = i - 1 \\ \int_0^\Delta [\langle S'_i(\omega, s) e_a, S'_i(\omega, s) e_c \rangle + \langle V'_{i+1}(\omega, s) e_a, V'_{i+1}(\omega, s) e_c \rangle] ds & \text{if } j = i \\ \int_0^\Delta \langle V'_{i+1}(\omega, s) e_a, S'_{i+1}(\omega, s) e_c \rangle ds & \text{if } j = i + 1 \\ 0 & \text{otherwise} \end{cases} \\
&= \begin{cases} \int_0^\Delta \langle e_a, S'_i(\omega, s)^T V'_i(\omega, s) e_c \rangle ds & \text{if } j = i - 1 \\ \int_0^\Delta [\langle e_a, S'_i(\omega, s)^T S'_i(\omega, s) e_c \rangle + \langle e_a, V'_{i+1}(\omega, s)^T V'_{i+1}(\omega, s) e_c \rangle] ds & \text{if } j = i \\ \int_0^\Delta \langle e_a, V'_{i+1}(\omega, s)^T S'_{i+1}(\omega, s) e_c \rangle ds & \text{if } j = i + 1 \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}$$

Let us define the block matrix function of  $s \in [0, \Delta]$  by,

$$\mathcal{F}_{ij}^n(\omega, s) = \delta_{ij} S'_i(\omega, s) + \delta_{i,j+1} V'_i(\omega, s)$$

or equivalently as

$$\mathcal{F}^n(\omega, s) := \begin{bmatrix} S'_1(\omega, s) & 0 & \dots & 0 & 0 \\ V'_2(\omega, s) & S'_2(\omega, s) & 0 & & 0 \\ 0 & V'_3(\omega, s) & S'_3(\omega, s) & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & V'_n(\omega, s) & S'_n(\omega, s) \end{bmatrix}. \quad (4.7)$$

where

$$V'_i(\omega, s) := C'_i(\omega, s) S_{i-1}(\omega, \Delta) - S'_i(\omega, s) F_{i-1}(\omega).$$

**Remark 4.6** *When  $\omega = 0$  we have  $S_i(0, s) = sI$ ,  $C_i(0, s) = I$ ,  $F_i(0) := \Delta^{-1} I \Delta = I$ ,  $V_i(0, s) := \Delta I - sI$ , and  $\mathcal{F}^n(0, s) = \mathcal{T}^n$  where  $\mathcal{T}_{ij}^n = (\delta_{ij} - \delta_{i,j+1}) I$ , i.e.*

$$\mathcal{F}^n(0, s) = \mathcal{T}^n := \begin{bmatrix} I & 0 & \dots & 0 & 0 \\ -I & I & 0 & \dots & 0 \\ 0 & -I & I & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & -I & I \end{bmatrix}. \quad (4.8)$$

It is also worth observing that

$$\mathcal{F}^n(\omega, 0) := \begin{bmatrix} I & 0 & \dots & 0 & 0 \\ -F_1(\omega) & I & 0 & \dots & 0 \\ 0 & -F_2(\omega) & I & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & -F_{n-1}(\omega) & I \end{bmatrix}, \quad (4.9)$$

or equivalently that

$$\mathcal{F}_{ki}^n(\omega, 0) = \delta_{ki}I - \delta_{i,k-1}F_i(\omega). \quad (4.10)$$

**Theorem 4.7** Let  $\mathcal{F}^n(s)_{ij} = \delta_{ij}S'_i(s) + \delta_{i,j+1}V'_i(s)$  (See Equation (4.7).) Then

$$\det \left( \left\{ \int_0^1 \langle f'_{i,a}(\omega, s), f'_{j,c}(\omega, s) \rangle ds \right\} \right) = \det \left( \int_0^\Delta (\mathcal{F}^n(\omega, s))^T \mathcal{F}^n(\omega, s) ds \right), \quad (4.11)$$

$$\det \left( \left\{ \sum_{k=1}^n \langle f'_{i,a}(\omega, s_{k-1}+), f'_{j,c}(\omega, s_{k-1}+) \rangle \Delta \right\} \right) = \Delta^{nd}, \quad (4.12)$$

and hence

$$(\rho_n \circ \phi)^2(\omega) = \det \left( \frac{1}{\Delta} \int_0^\Delta (\mathcal{F}^n(\omega, s))^T \mathcal{F}^n(\omega, s) ds \right). \quad (4.13)$$

**Proof.** Since

$$(\mathcal{F}_{ij}^n(\omega, s))^T = \delta_{ij}S'_i(\omega, s)^T + \delta_{j,i+1}V'_j(\omega, s)^T,$$

$$\begin{aligned} & [(\mathcal{F}^n(\omega, s))^T \mathcal{F}^n(\omega, s)]_{ij} \\ &= \sum_k \left[ \delta_{ik}S'_i(\omega, s)^T + \delta_{k,i+1}V'_k(\omega, s)^T \right] \left[ \delta_{kj}S'_k(\omega, s) + \delta_{k,j+1}V'_k(\omega, s) \right] \\ &= \delta_{ij}S'_i(\omega, s)^T S'_i(\omega, s) + \delta_{i,j+1}S'_i(\omega, s)^T V'_j(\omega, s) + \delta_{i+1,j}V'_{i+1}(\omega, s)^T S'_{i+1}(\omega, s) + \delta_{ij}V'_{i+1}(\omega, s)^T V'_{i+1}(\omega, s) \\ &= \delta_{ij} \left[ S'_i(\omega, s)^T S'_i(\omega, s) + V'_{i+1}(\omega, s)^T V'_{i+1}(\omega, s) \right] + \delta_{i,j+1}S'_i(\omega, s)^T V'_j(\omega, s) + \delta_{i+1,j}V'_{i+1}(\omega, s)^T S'_{i+1}(\omega, s). \end{aligned}$$

So comparing with the results from Lemma 4.5 it follows that

$$\begin{aligned} G^1(X^{f_{i,a}}, X^{f_{j,c}})(\omega) &= \int_0^1 g \left( \frac{\nabla X^{f_{i,a}}(\omega, s)}{ds}, \frac{\nabla X^{f_{j,c}}(\omega, s)}{ds} \right) ds \\ &= \int_0^1 \langle f'_{i,a}(\omega, s), f'_{j,c}(\omega, s) \rangle ds = \int_0^1 \left\langle e_a, [(\mathcal{F}^n(\omega, s))^T \mathcal{F}^n(\omega, s)]_{ij} e_c \right\rangle ds, \end{aligned}$$

from which Equation (4.11) follows.

In order to prove Equation (4.12) we begin by observing that

$$\begin{aligned}
f'_{i,a}(\omega, s_{k-1}+) &= [1_{J_i}(s_{k-1}+) S'_i(\omega, s_{k-1} - s_{i-1}) + 1_{J_{i+1}}(s_{k-1}+) V'_{i+1}(\omega, s_{k-1} - s_i)] e_a \\
&= [\delta_{ik} S'_i(\omega, 0) + \delta_{i,k-1} V'_{i+1}(\omega, 0)] e_a \\
&= [\delta_{ik} I - \delta_{i,k-1} F_i(\omega)] e_a = \mathcal{F}_{ik}^n(\omega, 0) e_a
\end{aligned}$$

where the last equality follows from Equation (4.10). Hence it follows that

$$\begin{aligned}
G_{\mathcal{P}}^1(X^{f_{i,a}}, X^{f_{j,c}})(\omega) &= \sum_{k=1}^n g\left(\frac{\nabla X^{f_{i,a}}(\omega, s_{k-1}+)}{ds}, \frac{\nabla X^{f_{j,c}}(\omega, s_{k-1}+)}{ds}\right) \\
&= \sum_{k=1}^n \langle f'_{i,a}(\omega, s_{k-1}+), f'_{j,c}(\omega, s_{k-1}+) \rangle \Delta = \Delta \sum_{k=1}^n \langle \mathcal{F}_{ki}^n(\omega, 0) e_a, \mathcal{F}_{kj}^n(\omega, 0) e_c \rangle \\
&= \Delta \sum_{k=1}^n \langle e_a, (\mathcal{F}_{ik}^n(\omega, 0))^T \mathcal{F}_{kj}^n(\omega, 0) e_c \rangle = \Delta \langle e_a, [(\mathcal{F}^n(\omega, 0))^T \mathcal{F}^n(\omega, 0)]_{ij} e_c \rangle
\end{aligned}$$

and therefore

$$\det\left(\left\{\sum_{k=1}^n f'_{i,a}(\omega, s_{k-1}+) \cdot f'_{j,c}(\omega, s_{k-1}+) \Delta\right\}\right) = \det(\Delta(\mathcal{F}^n(\omega, 0))^T \mathcal{F}^n(\omega, 0)) = \Delta^{nd}.$$

Equation (4.13) now follows from Equation (2.5). ■

## 4.1 Some Identities

**Definition 4.8** For real square matrix functions,  $A(s)$  and  $B(s)$ , of  $s \in [0, \Delta]$ , let

$$\langle A \rangle := \frac{1}{\Delta} \int_0^\Delta A(s) ds$$

and

$$\begin{aligned}
\text{Cov}(A, B) &= \frac{1}{\Delta} \int_0^\Delta A(s)^T B(s) ds - \left(\frac{1}{\Delta} \int_0^\Delta A(s) ds\right)^T \left(\frac{1}{\Delta} \int_0^\Delta B(s) ds\right) \\
&= \langle A^T B \rangle - \langle A \rangle^T \langle B \rangle.
\end{aligned}$$

Notice that  $\langle A \rangle$  and  $\text{Cov}(A, B)$  is again a square matrix.

The following proposition summarizes some basic and easily proved properties of  $\text{Cov}(A, B)$ .

**Proposition 4.9** The *covariance functional*,  $\text{Cov}$ , has the following properties:

1.  $\text{Cov}(A, B)$  is bilinear in  $A$  and  $B$ .

2.  $\text{Cov}(A, B)$  may be computed as

$$\begin{aligned}\text{Cov}(A, B) &= \frac{1}{\Delta} \int_0^\Delta [A(s) - \langle A \rangle]^T [B(s) - \langle B \rangle] ds \\ &= \left\langle [A(\cdot) - \langle A \rangle]^T [B(\cdot) - \langle B \rangle] \right\rangle.\end{aligned}$$

3.  $\text{Cov}(A, B) = 0$  if either  $A(s)$  or  $B(s)$  is a constant function.

4.  $\text{Cov}(A, A)$  is always a symmetric non-negative matrix.

**Note :** To simplify notation, for the rest of this section we will typically be omitting the argument,  $\omega$ , from the expressions to follow.

**Definition 4.10** Define  $\mathcal{G}^n(s) := \mathcal{F}^n(s) - \mathcal{T}^n = \mathcal{F}^n(s) - \mathcal{F}^n(0)$ , i.e.

$$\begin{aligned}\mathcal{G}_{ij}^n(s) &= \delta_{ij} [S'_i(s) - I] + \delta_{i,j+1} [V'_i(s) + I] \\ &= \delta_{ij} [S'_i(s) - I] + \delta_{i,j+1} [C'_i(s)S_{i-1}(\Delta) - S'_i(s)F_{i-1} + I].\end{aligned}\tag{4.14}$$

Also let

$$\mathcal{Y}^n := \langle \mathcal{G}^n \rangle = \frac{1}{\Delta} \int_0^\Delta \mathcal{G}^n(s) ds.\tag{4.15}$$

**Lemma 4.11** Let  $\mathcal{V}_{ij}^n = \delta_{ij} \frac{1}{\Delta} S_i(\Delta)$ , i.e.

$$\mathcal{V}^n := \frac{1}{\Delta} \begin{bmatrix} S_1(\Delta) & 0 & \dots & 0 \\ 0 & S_2(\Delta) & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & S_n(\Delta) \end{bmatrix}\tag{4.16}$$

and  $\mathcal{D}_{ij}^n := \delta_{ij} [\frac{1}{\Delta} S_i(\Delta) - I]$ , i.e.  $\mathcal{D}^n = \mathcal{V}^n - \mathcal{I}^n$ . Then  $\mathcal{Y}^n = \mathcal{T}^n \mathcal{D}^n$  and

$$\begin{aligned}\langle (\mathcal{F}^n)^T \mathcal{F}^n \rangle &= (\mathcal{T}^n + \mathcal{Y}^n)^T (\mathcal{T}^n + \mathcal{Y}^n) + \text{Cov}(\mathcal{G}^n, \mathcal{G}^n) \\ &= (\mathcal{V}^n)^T (\mathcal{T}^n)^T \mathcal{T}^n \mathcal{V}^n + \text{Cov}(\mathcal{G}^n, \mathcal{G}^n).\end{aligned}\tag{4.17}$$

**Proof.** By the fundamental theorem of calculus along with Equations (4.5) and (4.6) we have

$$\begin{aligned}\mathcal{Y}_{ij}^n &= \frac{1}{\Delta} \int_0^\Delta \left( \delta_{ij} [S'_i(s) - I] + \delta_{i,j+1} [V'_i(s) + I] \right) ds \\ &= \frac{1}{\Delta} (\delta_{ij} [S_i(\Delta) - I\Delta] + \delta_{i,j+1} [V_i(\Delta) - V_i(0) + \Delta I]) \\ &= \delta_{ij} \left[ \frac{S_i(\Delta)}{\Delta} - I \right] - \delta_{i,j+1} \left[ \frac{S_{i-1}(\Delta)}{\Delta} - I \right].\end{aligned}$$



On the other hand

$$\begin{aligned} (\mathcal{T}^n \mathcal{D}^n)_{ij} &= \sum_k [\delta_{ik} - \delta_{i,k+1}] I \delta_{kj} \left[ \frac{1}{\Delta} S_k(\Delta) - I \right] \\ &= \delta_{ij} \left[ \frac{1}{\Delta} S_i(\Delta) - I \right] - \delta_{i-1,j} \left[ \frac{1}{\Delta} S_{i-1}(\Delta) - I \right] = \mathcal{Y}_{ij}^n. \end{aligned}$$

The second assertion is a consequence of the following simple manipulations,

$$\begin{aligned} \langle (\mathcal{F}^n)^T \mathcal{F}^n \rangle &= \langle (\mathcal{T}^n + \mathcal{G}^n)^T (\mathcal{T}^n + \mathcal{G}^n) \rangle \\ &= (\mathcal{T}^n)^T \mathcal{T}^n + (\mathcal{T}^n)^T \mathcal{Y}^n + (\mathcal{Y}^n)^T \mathcal{T}^n + \langle (\mathcal{G}^n)^T \mathcal{G}^n \rangle \\ &= (\mathcal{T}^n + \mathcal{Y}^n)^T (\mathcal{T}^n + \mathcal{Y}^n) + \langle (\mathcal{G}^n)^T \mathcal{G}^n \rangle - (\mathcal{Y}^n)^T \mathcal{Y}^n \\ &= (\mathcal{T}^n + \mathcal{Y}^n)^T (\mathcal{T}^n + \mathcal{Y}^n) + \text{Cov}(\mathcal{G}^n, \mathcal{G}^n). \end{aligned}$$

This completes the proof since

$$\mathcal{T}^n + \mathcal{Y}^n = \mathcal{T}^n (\mathcal{I}^n + \mathcal{D}^n) = \mathcal{T}^n \mathcal{V}^n.$$

■

**Corollary 4.12** *Letting  $\mathcal{M}^n = \text{Cov}(\mathcal{G}^n, \mathcal{G}^n)$ ,*

$$\mathcal{S}^n := (\mathcal{T}^n)^{-1} = \begin{bmatrix} I & 0 & \dots & 0 \\ I & I & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ I & \dots & I & I \end{bmatrix}, \quad (4.18)$$

and  $\mathcal{V}^n$  be as in Equation (4.16), dropping the superscript  $n$ , we have

$$\langle \mathcal{F}^T \mathcal{F} \rangle = \mathcal{V}^T \mathcal{T}^T \left( \mathcal{I} + \mathcal{S}^T (\mathcal{V}^T)^{-1} \mathcal{M} \mathcal{V}^{-1} \mathcal{S} \right) \mathcal{T} \mathcal{V}$$

and

$$\det(\langle \mathcal{F}^T \mathcal{F} \rangle) = [\det(\mathcal{V})]^2 \cdot \det\left(\mathcal{I} + \mathcal{S}^T (\mathcal{V}^T)^{-1} \mathcal{M} \mathcal{V}^{-1} \mathcal{S}\right). \quad (4.19)$$

## 4.2 The Key Determinant Formula

Our next goal is to expand out  $\mathcal{V}^n$  and  $((\mathcal{V}^n)^T)^{-1} \mathcal{M}^n (\mathcal{V}^n)^{-1}$  in powers of  $\omega$ . It turns out that we need the expansion of  $\mathcal{V}^n$  to order  $\|\omega\|^3$  and  $((\mathcal{V}^n)^T)^{-1} \mathcal{M}^n (\mathcal{V}^n)^{-1}$  to order  $\|\omega\|^5$ .

**Notation 4.13** *Recall the definition of  $A_i(\omega, s)$  in Equation (3.4). We will write  $A_i(0) = A_i(\omega, 0)$  and also  $A_i(t) = A_i(\omega, t)$  to simplify the notation. And when we write  $y = O(x)$ , we mean there exists some constant  $C > 0$  independent of  $i, n$  and  $\omega$  such that  $\|y\| \leq C \|x\|$ .*

**Proposition 4.14** *There exists a  $C < \infty$  such that the following estimates hold for  $0 \leq s \leq \Delta$ ;*

$$\left\| S'_i(s) - \left( I + \frac{1}{2} A_i(0) s^2 \right) \right\| \leq C \|\Delta_i \omega\|^3, \quad (4.20)$$

$$\left\| \frac{S_i(s)}{s} - \left( I + \frac{1}{6} A_i(0) s^2 \right) \right\| \leq C \|\Delta_i \omega\|^3, \quad (4.21)$$

$$\left\| C_i(s) - \left( I + \frac{1}{2} A_i(0) s^2 \right) \right\| \leq C \|\Delta_i \omega\|^3 \text{ and} \quad (4.22)$$

$$\|C'_i(s) - A_i(0) s\| \leq C s^{-1} \|\Delta_i \omega\|^3. \quad (4.23)$$

In the sequel we will abbreviate these type of estimates by writing

$$S'_i(s) = I + \frac{1}{2} A_i(0) s^2 + O\left(\|\Delta_i \omega\|^3\right), \quad (4.24)$$

$$\frac{S_i(s)}{s} = I + \frac{1}{6} A_i(0) s^2 + O\left(\|\Delta_i \omega\|^3\right), \quad (4.25)$$

$$C_i(s) = I + \frac{1}{2} A_i(0) s^2 + O\left(\|\Delta_i \omega\|^3\right) \text{ and} \quad (4.26)$$

$$C'_i(s) = s^{-1} \left[ A_i(0) s^2 + O\left(\|\Delta_i \omega\|^3\right) \right]. \quad (4.27)$$

**Proof.** Let  $(DA_i)(\omega'(s), \cdot, \cdot) := \frac{d}{ds} \Omega_{u(s_{i-1}+s)}$ . Since  $M$  is compact, there exists a constant  $C > 0$  such that  $\sup_{u \in O(M)} \|D\Omega_u\| < C$ . Then

$$\begin{aligned} \|(DA_i)(\omega'(s), \omega'(s), \cdot) \omega'(s)\| &\leq \frac{1}{\Delta^3} (\|(DA_i)(\Delta_i \omega, \Delta_i \omega, \cdot) \Delta_i \omega\|) \\ &= O\left(\frac{1}{\Delta^3} \|\Delta_i \omega\|^3\right). \end{aligned}$$

Thus

$$\int_0^s r A_i(r) dr = \int_0^s r \left[ A_i(0) + \int_0^r A'_i(t) dt \right] dr = \frac{1}{2} A_i(0) s^2 + O\left(\|\Delta_i \omega\|^3\right)$$

and similarly that

$$\begin{aligned} \frac{1}{s} \int_0^s (s-r) r A_i(r) dr &= \frac{1}{s} \int_0^s (s-r) r A_i(0) dr + O\left(\|\Delta_i \omega\|^3\right) \\ &= \frac{1}{6} A_i(0) s^2 + O\left(\|\Delta_i \omega\|^3\right) \end{aligned}$$

and finally

$$\int_0^s (s-u) A_i(u) du = \int_0^s (s-u) A_i(0) du + O\left(\|\Delta_i \omega\|^3\right) = \frac{1}{2} A_i(0) s^2 + O\left(\|\Delta_i \omega\|^3\right).$$

Combining these results with the three estimates in Proposition 3.12 with  $\kappa := K \|\omega'_i\|^2$  proves (4.20) – (4.22). For Equation (4.23), we have

$$\begin{aligned} C'_i(s) &= \int_0^s A_i(r) C_i(r) dr = \int_0^s \left[ A_i(0) + \int_0^r A'_i(t) dt \right] \left( I + O\left(\|\Delta_i \omega\|^2\right) \right) dr \\ &= A_i(0) s + s^{-1} O\left(\|\Delta_i \omega\|^3\right) \end{aligned}$$

as desired. ■

**Corollary 4.15** With  $\mathcal{V}^n = \mathcal{I}^n + \mathcal{D}^n$  as in Equation (4.16) we have

$$\mathcal{V}_{ij}^n = \delta_{ij} \left( I + \frac{1}{6} A_i(0) \Delta^2 + \eta_i(\Delta) \right) \quad (4.28)$$

where

$$\eta_i(s) := \frac{S_i(s)}{s} - \left( I + \frac{1}{6} A_i(0) s^2 \right) \quad (4.29)$$

and  $\|\eta_i(\Delta)\| = O(\|\Delta_i \omega\|^3)$ .

**Lemma 4.16** The function,  $F_i$  in Equation (4.4) satisfies,

$$F_i = I + \frac{1}{6} A_i(0) \Delta^2 + \frac{1}{3} A_{i+1}(0) \Delta^2 + O(\|\Delta_i \omega\|^3 \vee \|\Delta_{i+1} \omega\|^3). \quad (4.30)$$

**Proof.** In order to simplify notation, let  $a_i := A_i(0) \Delta^2$  and  $\beta_i = \|\Delta_i \omega\|^3$ . Then

$$\begin{aligned} F_i &= S_{i+1}(\Delta)^{-1} C_{i+1}(\Delta) S_i(\Delta) = \left( \frac{S_{i+1}(\Delta)}{\Delta} \right)^{-1} C_{i+1}(\Delta) \frac{S_i(\Delta)}{\Delta} \\ &= \left( I + \frac{1}{6} a_{i+1} + O(\beta_{i+1}) \right)^{-1} \left( I + \frac{1}{2} a_{i+1} + O(\beta_{i+1}) \right) \left( I + \frac{1}{6} a_i + O(\beta_i) \right) \\ &= \left( I - \frac{1}{6} a_{i+1} + O(\beta_{i+1}) \right) \left( I + \frac{1}{2} a_{i+1} + O(\beta_{i+1}) \right) \left( I + \frac{1}{6} a_i + O(\beta_i) \right) \\ &= I + \left( \frac{1}{2} - \frac{1}{6} \right) a_{i+1} + \frac{1}{6} a_i + O(\|\Delta_i \omega\|^3) + O(\|\Delta_{i+1} \omega\|^3) \end{aligned}$$

which is equivalent to Equation (4.30). ■

**Theorem 4.17** Let

$$\mathcal{H}_{ij}^n(s) := \delta_{ij} A_i(0) \frac{s^2}{2} + \delta_{i,j+1} \left[ A_i(0) \left( s\Delta - \frac{s^2}{2} - \frac{\Delta^2}{3} \right) - A_{i-1}(0) \frac{\Delta^2}{6} \right]$$

and  $\Upsilon_{ij}^n(s) := \mathcal{G}_{ij}^n(s) - \mathcal{H}_{ij}^n(s)$ . Then  $\|\Upsilon_{ij}^n(s)\| = O(\|\Delta_{i-1} \omega\|^3 \vee \|\Delta_i \omega\|^3)$  and since  $\Upsilon_{ij}^n = 0$  unless  $i \in \{j, j+1\}$ , it follows that

$$\mathcal{G}^n(s) = \mathcal{H}^n(s) + \Upsilon^n(s) \text{ and } \|\Upsilon^n\| = O\left(\bigvee_{i=1,\dots,n} \|\Delta_i \omega\|^3\right). \quad (4.31)$$

**Proof.** Let  $\gamma_i = \|\Delta_{i-1} \omega\|^3 \vee \|\Delta_i \omega\|^3$ . By Proposition 4.14 and Lemma 4.16,

$$\begin{aligned} C'_i(s) S_{i-1}(\Delta) &= \left[ A_i(0) s\Delta + O(\|\Delta_i \omega\|^3) \right] \Delta^{-1} S_{i-1}(\Delta) \\ &= \left[ A_i(0) s\Delta + O(\|\Delta_i \omega\|^3) \right] \left[ I + \frac{1}{6} A_{i-1}(0) \Delta^2 + O(\|\Delta_{i-1} \omega\|^3) \right] \\ &= A_i(0) s\Delta + O(\gamma_i) \end{aligned} \quad (4.32)$$

and

$$\begin{aligned}
& S'_i(s)F_{i-1} \\
&= \left( I + \frac{1}{2}A_i(0)s^2 + O(\|\Delta_i\omega\|^3) \right) \left( I + \frac{1}{6}A_{i-1}(0)\Delta^2 + \frac{1}{3}A_i(0)\Delta^2 + O(\gamma_i) \right) \\
&= I + A_{i-1}(0)\frac{\Delta^2}{6} + A_i(0)\left(\frac{s^2}{2} + \frac{\Delta^2}{3}\right) + O(\gamma_i).
\end{aligned} \tag{4.33}$$

Combining the last two equations shows,

$$\begin{aligned}
V'_i(s) &= C'_i(s)S_{i-1}(\Delta) + I - S'_i(s)F_{i-1} \\
&= A_i(0)\left(s\Delta - \frac{s^2}{2} - \frac{\Delta^2}{3}\right) - A_{i-1}(0)\frac{\Delta^2}{6} + O(\|\Delta_{i-1}\omega\|^3 \vee \|\Delta_i\omega\|^3).
\end{aligned} \tag{4.34}$$

This equation along with Equations (4.14) and (4.24) shows

$$\begin{aligned}
\mathcal{G}_{ij}^n(s) &= \delta_{ij} \left[ S'_i(s) - I \right] + \delta_{i,j+1} [V'_i(s) + I] \\
&= \delta_{ij} \left[ A_i(0)\frac{s^2}{2} + O(\|\Delta_i\omega\|^3) \right] \\
&\quad + \delta_{i,j+1} \left[ A_i(0)\left(s\Delta - \frac{s^2}{2} - \frac{\Delta^2}{3}\right) - A_{i-1}(0)\frac{\Delta^2}{6} + O(\|\Delta_{i-1}\omega\|^3 \vee \|\Delta_i\omega\|^3) \right] \\
&= \mathcal{H}_{ij}^n(s) + O(\|\Delta_{i-1}\omega\|^3 \vee \|\Delta_i\omega\|^3).
\end{aligned}$$

■

**Theorem 4.18** *The matrix  $\mathcal{M}^n$  of Corollary 4.12 satisfies,  $\mathcal{M}^n = \mathcal{C}^n + \tilde{\mathcal{E}}^n$  where  $\tilde{\mathcal{E}}^n$  is a tri-block-diagonal matrix such that  $\|\tilde{\mathcal{E}}^n\| = O(\bigvee_{i=1,\dots,n} \|\Delta_i\omega\|^5)$  and  $\mathcal{C}^n$  is the non-negative tri-block-diagonal matrix given by*

$$\mathcal{C}_{ij}^n := \delta_{ij} \frac{\Delta^4}{45} [A_i^2(0) + A_{i+1}^2(0)] + \frac{7}{360} [\delta_{i,j+1} A_i^2(0) + \delta_{i,j-1} A_j^2(0)] \Delta^4, \tag{4.35}$$

where  $A_{n+1}^2(0) := 0$ . Equivalently,

$$\mathcal{C}^n := \begin{pmatrix} \frac{1}{45}(a_1^2 + a_2^2) & \frac{7}{360}a_2^2 & 0 & \cdots & 0 \\ \frac{7}{360}a_2^2 & \frac{1}{45}(a_2^2 + a_3^2) & \frac{7}{360}a_3^2 & \ddots & \vdots \\ 0 & \frac{7}{360}a_3^2 & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \frac{1}{45}(a_{n-1}^2 + a_n^2) & \frac{7}{360}a_n^2 \\ 0 & \cdots & 0 & \frac{7}{360}a_n^2 & \frac{1}{45}a_n^2 \end{pmatrix}$$

where as before,  $a_i := A_i(0)\Delta^2$ .

**Proof.** Since  $\mathcal{G}^n = \mathcal{H}^n + \Upsilon^n$  with  $\|\mathcal{H}^n\| = O\left(\bigvee_{i=1,\dots,n} \|\Delta_i \omega\|^2\right)$  and  $\|\Upsilon^n\| = O\left(\bigvee_{i=1,\dots,n} \|\Delta_i \omega\|^3\right)$ , it follows that

$$\begin{aligned}\mathcal{M}^n &= \text{Cov}(\mathcal{G}^n, \mathcal{G}^n) = \text{Cov}(\mathcal{H}^n + \Upsilon^n, \mathcal{H}^n + \Upsilon^n) \\ &= \text{Cov}(\mathcal{H}^n, \mathcal{H}^n) + \text{Cov}(\mathcal{H}^n, \Upsilon^n) + \text{Cov}(\Upsilon^n, \mathcal{H}^n) + \text{Cov}(\Upsilon^n, \Upsilon^n) \\ &= \text{Cov}(\mathcal{H}^n, \mathcal{H}^n) + \tilde{\mathcal{E}}^n = \mathcal{C}^n + \tilde{\mathcal{E}}^n\end{aligned}$$

where  $\mathcal{C}^n := \text{Cov}(\mathcal{H}^n, \mathcal{H}^n)$  and

$$\tilde{\mathcal{E}}^n = \text{Cov}(\mathcal{H}^n, \Upsilon^n) + \text{Cov}(\Upsilon^n, \mathcal{H}^n) + \text{Cov}(\Upsilon^n, \Upsilon^n) = O\left(\bigvee_{i=1,\dots,n} \|\Delta_i \omega\|^5\right).$$

Since  $\mathcal{H}^n = \mathcal{K}^n + \mathcal{L}^n$  where

$$\mathcal{K}_{ij}^n(s) := \delta_{ij} A_i(0) \frac{s^2}{2} + \delta_{i,j+1} A_i(0) \left[ s\Delta - \frac{s^2}{2} \right]$$

and  $\mathcal{L}^n$  is the constant (in  $s$ ) matrix,

$$\mathcal{L}_{ij}^n := -\delta_{i,j+1} \left[ A_i(0) \frac{\Delta^2}{3} + A_{i-1}(0) \frac{\Delta^2}{6} \right],$$

it follows, using Proposition 4.9, that

$$\mathcal{C}^n = \text{Cov}(\mathcal{H}^n, \mathcal{H}^n) = \text{Cov}(\mathcal{K}^n, \mathcal{K}^n).$$

Since

$$\langle \mathcal{K}_{ij}^n \rangle := \left[ \frac{1}{6} \delta_{ij} + \frac{1}{3} \delta_{i,j+1} \right] A_i(0) \Delta^2,$$

we have

$$\begin{aligned}\hat{\mathcal{K}}_{ij}^n(s) &:= \mathcal{K}_{ij}^n(s) - \langle \mathcal{K}_{ij}^n \rangle \\ &= \delta_{ij} A_i(0) \left( \frac{s^2}{2} - \frac{\Delta^2}{6} \right) + \delta_{i,j+1} A_i(0) \left( s\Delta - \frac{s^2}{2} - \frac{1}{3} \Delta^2 \right).\end{aligned}$$

Let

$$\alpha = \frac{s^2}{2} - \frac{\Delta^2}{6}, \quad \beta = s\Delta - \frac{s^2}{2} - \frac{1}{3} \Delta^2.$$

By direct integration, we get

$$\begin{aligned}\langle \alpha^2 \rangle &= \frac{1}{\Delta} \int_0^\Delta \alpha^2 ds = \frac{1}{\Delta} \int_0^\Delta \left( \frac{s^2}{2} - \frac{\Delta^2}{6} \right)^2 ds = \frac{\Delta^4}{45}, \\ \langle \beta^2 \rangle &= \frac{1}{\Delta} \int_0^\Delta \beta^2 ds = \frac{1}{\Delta} \int_0^\Delta \left( s\Delta - \frac{s^2}{2} - \frac{1}{3} \Delta^2 \right)^2 ds = \frac{\Delta^4}{45} \text{ and} \\ \langle \alpha\beta \rangle &= \frac{1}{\Delta} \int_0^\Delta \alpha\beta ds = \frac{1}{\Delta} \int_0^\Delta \left( \frac{s^2}{2} - \frac{\Delta^2}{6} \right) \left( s\Delta - \frac{s^2}{2} - \frac{1}{3} \Delta^2 \right) ds = \frac{7\Delta^4}{360}.\end{aligned}$$

Then we may conclude that

$$\begin{aligned}
\mathcal{C}_{ij}^n &= \text{Cov}(\mathcal{K}^n, \mathcal{K}^n)_{ij} = \sum_k \left\langle \left( (\hat{\mathcal{K}}^n)^T \right)_{ik} \hat{\mathcal{K}}_{kj}^n \right\rangle = \sum_k \left\langle \left( \hat{\mathcal{K}}_{ki}^n \right)^T \hat{\mathcal{K}}_{kj}^n \right\rangle \\
&= \sum_k \left\langle [\alpha \delta_{ki} + \beta \delta_{k,i+1}] A_k(0) [\alpha \delta_{kj} + \beta \delta_{k,j+1}] A_k(0) \right\rangle \\
&= \frac{\Delta^4}{45} \sum_k [\delta_{ki} \delta_{kj} + \delta_{k,i+1} \delta_{k,j+1}] A_k^2(0) + \Delta^4 \frac{7}{360} \sum_k [\delta_{ki} \delta_{k,j+1} + \delta_{k,i+1} \delta_{kj}] A_k^2(0) \\
&= \frac{\Delta^4}{45} \delta_{ij} (A_i^2(0) + A_{i+1}^2(0)) + \Delta^4 \frac{7}{360} [\delta_{i,j+1} + \delta_{i+1,j}] A_i^2(0)
\end{aligned}$$

which is equivalent to Equation (4.35). ■

Putting together the previous estimates leads to the following key determinant formula.

**Theorem 4.19** *As above, let  $\mathcal{S}^n = (\mathcal{T}^n)^{-1}$ ,*

$$\begin{aligned}
\mathcal{V}_{ij}^n &= \delta_{ij} + \mathcal{D}_{ij}^n = \delta_{ij} \frac{1}{\Delta} S_i(\Delta) = \delta_{ij} \left( I + \frac{1}{6} A_i(0) \Delta^2 + \eta_i(\Delta) \right), \\
\mathcal{C}_{ij}^n &= \frac{\Delta^4}{45} \delta_{ij} [A_i^2(0) + A_{i+1}^2(0)] + \frac{7}{360} [\delta_{i,j+1} A_i^2(0) + \delta_{i,j-1} A_j^2(0)] \Delta^4
\end{aligned}$$

where  $\|\eta_i(\Delta)\| \leq C \|\Delta_i \omega\|^3$ ,  $C$  is independent of  $i, n, \omega$  and define

$$\mathcal{U}^n := (\mathcal{S}^n)^T \mathcal{C}^n \mathcal{S}^n. \quad (4.36)$$

Then there exists a tri-block-diagonal matrix  $\mathcal{E}^n = \mathcal{E}^n(\omega)$  such that

$$\|\mathcal{E}^n\| = O \left( \bigvee_{i=1, \dots, n} \|\Delta_i \omega\|^5 \right)$$

and

$$\det \left( \langle (\mathcal{F}^n)^T \mathcal{F} \rangle \right) = [\det(\mathcal{V}^n)]^2 \cdot \det(\mathcal{I}^n + \mathcal{U}^n) \cdot \det(\mathcal{I}^n + \mathcal{X}^n), \quad (4.37)$$

where  $\mathcal{X}^n := (\mathcal{I}^n + \mathcal{U}^n)^{-1} (\mathcal{S}^n)^T \mathcal{E}^n \mathcal{S}^n$ .

**Proof.** To ease the notation, we will drop the superscript  $n$  in this proof. From Equation (4.19) of Corollary 4.12, Corollary 4.15, and Theorem 4.18,

$$\det \left( \langle \mathcal{F}^T \mathcal{F} \rangle \right) = [\det(\mathcal{V})]^2 \cdot \det \left( \mathcal{I} + \mathcal{S}^T (\mathcal{V}^T)^{-1} (\mathcal{C} + \tilde{\mathcal{E}}) \mathcal{V}^{-1} \mathcal{S} \right). \quad (4.38)$$

Now write  $\mathcal{V}^{-1} = \mathcal{I} + \Psi$ , where

$$\Psi := \sum_{n=1}^{\infty} (-1)^n \mathcal{D}^n$$

is a block-diagonal matrix satisfying,  $\|\Psi\| = O\left(\bigvee_{i=1,\dots,n} \|\Delta_i \omega\|^2\right)$ . Hence for  $\epsilon > 0$  sufficiently small, if  $\bigvee_{i=1,\dots,n} \|\Delta_i \omega\|^2 \leq \epsilon$ , then  $\|\mathcal{V}^{-1}\| \leq 2$ . Furthermore,

$$\begin{aligned} (\mathcal{V}^T)^{-1} (\mathcal{C} + \tilde{\mathcal{E}}) \mathcal{V}^{-1} &= (\mathcal{V}^T)^{-1} \mathcal{C} \mathcal{V}^{-1} + (\mathcal{V}^T)^{-1} \tilde{\mathcal{E}} \mathcal{V}^{-1} \\ &= (\mathcal{I} + \Psi^T) \mathcal{C} (\mathcal{I} + \Psi) + (\mathcal{V}^T)^{-1} \tilde{\mathcal{E}} \mathcal{V}^{-1} = \mathcal{C} + \mathcal{E} \end{aligned}$$

where  $\mathcal{E}$  is the tri-block-diagonal matrix defined by

$$\mathcal{E} = \mathcal{C} \Psi + \Psi^T \mathcal{C} + \Psi^T \mathcal{C} \Psi + (\mathcal{V}^T)^{-1} \tilde{\mathcal{E}} \mathcal{V}^{-1}$$

and  $\mathcal{E}$  satisfies the norm estimate,  $\|\mathcal{E}\| = O\left(\bigvee_{i=1,\dots,n} \|\Delta_i \omega\|^5\right)$ . Putting these results back into Equation (4.38) shows

$$\begin{aligned} \det(\langle \mathcal{F}^T \mathcal{F} \rangle) &= [\det(\mathcal{V})]^2 \cdot \det(\mathcal{I} + \mathcal{S}^T [\mathcal{C} + \mathcal{E}] \mathcal{S}) \\ &= [\det(\mathcal{V})]^2 \cdot \det(\mathcal{I} + \mathcal{U} + \mathcal{S}^T \mathcal{E} \mathcal{S}) \\ &= [\det(\mathcal{V})]^2 \cdot \det\left((\mathcal{I} + \mathcal{U}) \left(\mathcal{I} + (\mathcal{I} + \mathcal{U})^{-1} \mathcal{S}^T \mathcal{E} \mathcal{S}\right)\right) \end{aligned}$$

from which the desired result follows. ■

## 5 Convergence of $\{\rho_n \circ \phi \circ b_n\}_{n=1}^\infty$ in $\mu$ -measure

Recall that  $H_{\mathcal{P}}^\epsilon(\mathbb{R}^d)$  was defined as

$$H_{\mathcal{P}}^\epsilon(\mathbb{R}^d) = \{\omega \in H_{\mathcal{P}}(\mathbb{R}^d) : \|\Delta_i \omega\| < \epsilon \ \forall i\}$$

where  $\Delta_i \omega = \omega(s_i) - \omega(s_{i-1})$ . Note that  $\mathcal{V}^n$  and  $\mathcal{X}^n$  are only defined on  $H_{\mathcal{P}}^\epsilon(\mathbb{R}^d)$  for some  $\epsilon$  satisfying Lemma 4.3.

**Notation 5.1** *By abuse of notation, we will now write  $A_i(\omega, 0) = A_i(\omega)$  from now on. This should not be confused with the notation  $A_i(t)$  as described in Notation 4.13, where  $t \in [0, \frac{1}{n})$  and we suppressed the argument  $\omega$ .*

Unless stated otherwise, we will only consider equally spaced partitions  $\mathcal{P}_n = \{0 < \frac{1}{n} < \dots < \frac{n}{n} = 1\}$ . By Theorem 4.7 and 4.19,  $\rho_n$  has been written as a product of 3 terms, namely  $[\det(\mathcal{V}^n)]^2$ ,  $\det(\mathcal{I}^n + \mathcal{U}^n)$  and  $\det(\mathcal{I}^n + \mathcal{X}^n)$  on  $H_{\mathcal{P}_n}^\epsilon(\mathbb{R}^d)$ . We will now show that the determinant given in Theorem 4.19 has a limit as  $|\mathcal{P}_n| \rightarrow 0$ . The limit for each term will be computed in this order.

The following theorem is the Wong-Zakai type approximation theorem for solutions to Stratonovich stochastic differential equations. This theorem is a special case of Theorem 5.7.3 and Example 5.7.4

in [Kun97]. Theorems of this type have a long history starting with Wong and Zakai [WZ65, WZ67]. The following version maybe found in [DH98].

**Theorem 5.2** *Let  $f : \mathbb{R}^d \times \mathbb{R}^n \rightarrow \text{End}(\mathbb{R}^d, \mathbb{R}^n)$  and  $f_0 : \mathbb{R}^d \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  be twice differentiable with bounded continuous derivatives. Let  $\xi_0 \in \mathbb{R}^n$  and  $\mathcal{P}$  be any partition of  $[0, 1]$ . Further, let  $b$  and  $b_{\mathcal{P}}$  be as in Definition 2.8 and  $\xi_{\mathcal{P}}(s)$  denote the solution to the ordinary differential equation*

$$\xi'_{\mathcal{P}}(s) = f(\xi_{\mathcal{P}}(s))b'_{\mathcal{P}}(s) + f_0(\xi_{\mathcal{P}}(s)), \quad \xi_{\mathcal{P}}(0) = \xi_0$$

*and  $\xi$  denote the solution to the Stratonovich stochastic differential equation,*

$$d\xi(s) = f(\xi(s))\delta b(s) + f_0(\xi(s))ds, \quad \xi(0) = \xi_0.$$

*Then for any  $\alpha \in (0, \frac{1}{2})$ ,  $p \in [1, \infty)$ , there is a constant  $C(p, \alpha) < \infty$  depending only on  $f, f_0$  and  $M$  so that*

$$\mathbb{E}[\sup_{s \leq 1} \|\xi_{\mathcal{P}}(s) - \xi(s)\|^p] \leq C(p, \alpha)|\mathcal{P}|^{\alpha p}.$$

**Definition 5.3** 1. *Let  $u$  be the solution to the Stratonovich stochastic differential equation*

$$\delta u = \mathcal{H}_u u \delta b, \quad u(0) = u_0. \quad (5.1)$$

*Hence  $u$  maybe viewed as  $\mu$ -a.s. defined function from  $W(\mathbb{R}^d) \rightarrow W(O(M))$ .*

2. *Let  $\tilde{\phi} := \pi \circ u : W(\mathbb{R}^d) \rightarrow W(M)$ . This map will be called the stochastic development map.*

The following facts will be stated without any proof. See [Dri92].

**Remark 5.4** 1. *Using Theorem 5.2, one may show that  $\tilde{\phi}$  is a "stochastic extension" of  $\phi$ , i.e.,  $\tilde{\phi} = \lim_{|\mathcal{P}| \rightarrow 0} \phi \circ b_{\mathcal{P}}$ . Moreover, the law of  $\tilde{\phi}$  (i.e.,  $\mu\tilde{\phi}^{-1}$ ) is the Wiener measure  $\nu$  on  $W(M)$ .*

2. *One can prove that  $u_s = \widetilde{//}_s(\tilde{\phi})$  where  $\widetilde{//}$  is stochastic parallel transport defined in Definition 1.12.*

3. *The law of  $u$  under  $\mu$  on  $W(\mathbb{R}^d)$  and the law of  $\widetilde{//}$  under  $\nu$  on  $W(M)$  are equal.*

**Definition 5.5** *Let  $0 < \alpha < \frac{1}{2}$  and define for  $\mathcal{P}_n = \{0 = \frac{1}{n} < \dots < \frac{n}{n} = 1\}$ ,*

$$W_{\alpha}(\mathbb{R}^d) = \left\{ \omega \in W(\mathbb{R}^d) \mid \exists n(\omega) \text{ s.t. } \forall n \geq n(\omega), \quad \bigvee_{i=1, \dots, n} \|\Delta_i \omega\| \leq n^{-\alpha} < \epsilon \right\}$$

*for some  $\epsilon$  satisfying Lemma 4.3 and  $n_0 : W_{\alpha}(\mathbb{R}^d) \rightarrow \mathbb{N}$  by*

$$n_0(\omega) := \inf \left\{ n(\omega) \in \mathbb{N} \mid \forall n \geq n(\omega), \quad \bigvee_{i=1, \dots, n} \|\Delta_i \omega\| \leq n^{-\alpha} < \epsilon \right\}.$$



Note that

$$(W_\alpha(\mathbb{R}^d))^c = \left\{ \omega \in W(\mathbb{R}^d) \mid \bigvee_{i=1, \dots, n} \|\Delta_i \omega\| > n^{-\alpha} \text{ i.o.} \right\}.$$

**Lemma 5.6** *Let  $0 < \alpha < \frac{1}{2}$ , then*

$$\mu(W_\alpha(\mathbb{R}^d)) = 1.$$

**Proof.** Now

$$\begin{aligned} \mu \left( \left\{ \omega \in W(\mathbb{R}^d) \mid \bigvee_{i=1, \dots, n} \|\Delta_i \omega\| > n^{-\alpha} \right\} \right) &\leq \frac{1}{n^{-\alpha p}} \mathbb{E} \left[ \bigvee_{i=1, \dots, n} \|\Delta_i b\|^p \right] \\ &\leq \frac{1}{n^{-\alpha p}} \mathbb{E} \left[ \sum_{i=1}^n \|\Delta_i b\|^p \right] = \frac{C_p}{n^{-\alpha p}} \left( \frac{1}{n} \right)^{\frac{p}{2}} n, \end{aligned}$$

where  $C_p > 0$  is some constant. Therefore,

$$\begin{aligned} \sum_{n=1}^{\infty} \mu \left( \left\{ \omega \in W(\mathbb{R}^d) \mid \bigvee_{i=1, \dots, n} \|\Delta_i \omega\| > n^{-\alpha} \right\} \right) \\ \leq C_p \sum_{n=1}^{\infty} \left( \frac{1}{n} \right)^{\frac{p}{2} - 1 - \alpha p} = C_p \sum_{n=1}^{\infty} \left( \frac{1}{n} \right)^{(\frac{1}{2} - \alpha)p - 1} < \infty, \end{aligned}$$

if  $p$  is sufficiently large. Hence by Borel Cantelli Lemma,

$$\mu \left( \left\{ \omega \in W(\mathbb{R}^d) \mid \bigvee_{i=1, \dots, n} \|\Delta_i \omega\| > n^{-\alpha} \text{ i.o.} \right\} \right) = 0,$$

and hence the proof. ■

Using Lemma 5.6, we can now extend the definition of  $\det(\mathcal{V}^n)$  and  $\det(\mathcal{I}^n + \mathcal{U}^n)$  to be a  $\mu$ -a.s. function. Let  $\omega \in W_\alpha(\mathbb{R}^d)$ . By Definition 5.5, for  $n \geq n_0(\omega)$ ,

$$\bigvee_{i=1, \dots, n} \|\Delta_i \omega\| < n^{-\alpha} < \epsilon.$$

Therefore  $b_n(\omega) \in H_{\mathcal{P}_n}^\epsilon(\mathbb{R}^d)$  for all  $n \geq n_0(\omega)$ .

**Definition 5.7** *Define*

$$\det(\widetilde{\mathcal{V}^n \circ b_n}) = 1_{\{n \geq n_0\}} \cdot \det(\mathcal{V}^n \circ b_n)$$

and

$$\det(\widetilde{\mathcal{I}^n + \mathcal{X}^n}) \circ b_n = 1_{\{n \geq n_0\}} \cdot \det(\mathcal{I}^n + \mathcal{X}^n) \circ b_n.$$

**Notation 5.8** *Throughout the next few sections, let  $b(s) : W(\mathbb{R}^d) \rightarrow \mathbb{R}^d$  be the projection map,  $b(s)(\omega) = \omega(s)$  for all  $0 \leq s \leq 1$  and  $\omega \in W(\mathbb{R}^d)$ . Note that when  $W(\mathbb{R}^d)$  is equipped with Wiener measure,  $\mu$ ,  $\{b(s) : 0 \leq s \leq 1\}$  is a Brownian motion. We further let  $\phi_n = \phi \circ b_n$  and  $u_n = //(\phi_n)$ .*

## 5.1 Convergence of $\det(\widetilde{\mathcal{V}^n \circ b_n})$

**Lemma 5.9**

$$\sum_{i=1}^n \left( \operatorname{tr} A_i(b_n) \Delta_i s^2 - \left( - \sum_{i=1}^n \operatorname{Scal}(\phi_n(s_{i-1})) \frac{1}{n} \right) \right) \longrightarrow 0$$

$\mu - a.s.$  as  $n \rightarrow \infty$ .

**Proof.** Note that

$$\Delta_i b_n = b_n(s_i) - b_n(s_{i-1}) = b(s_i) - b(s_{i-1}) = \Delta_i b$$

and

$$A_i(b_n) \Delta_i s^2 = \Omega_{u_n(s_{i-1})}((b_n)'_i, \cdot) (b_n)'_i \Delta_i s^2 = \Omega_{u_n(s_{i-1})}(\Delta_i b_n, \cdot) \Delta_i b_n.$$

Let  $Ric_{u_n(s)} := \sum_{i=1}^d \Omega_{u_n(s)}(\cdot, e_i) e_i$ . Using the symmetry of  $Ric$ ,

$$\begin{aligned} \sum_{i=1}^n \operatorname{tr} A_i(b_n) \Delta_i s^2 &= \sum_{i=1}^n \operatorname{tr} \Omega_{u_n(s_{i-1})}(\Delta_i b_n, \cdot) \Delta_i b_n = - \sum_{i=1}^n \left\langle Ric_{u_n(s_{i-1})} \Delta_i b_n, \Delta_i b_n \right\rangle \\ &= - \sum_{i=1}^n \left\langle Ric_{u_n(s_{i-1})} \Delta_i b, \Delta_i b \right\rangle. \end{aligned}$$

By Ito's formula,

$$\begin{aligned} &\left\langle Ric_{u_n(s_{i-1})} \Delta_i b, \Delta_i b \right\rangle \\ &= \left\langle Ric_{u_n(s_{i-1})}(b(s_i) - b(s_{i-1})), b(s_i) - b(s_{i-1}) \right\rangle \\ &= 2 \int_{s_{i-1}}^{s_i} \left\langle Ric_{u_n(s_{i-1})}((b(s) - b(s_{i-1}))), db(s) \right\rangle + \int_{s_{i-1}}^{s_i} \operatorname{tr} Ric_{u_n(s_{i-1})} ds \\ &= 2 \int_{s_{i-1}}^{s_i} \left\langle Ric_{u_n(s_{i-1})}(b(s) - b(s_{i-1})), db(s) \right\rangle + \operatorname{Scal}(\phi_n(s_{i-1})) \Delta_i s. \end{aligned}$$

Thus

$$\sum_{i=1}^n \left( \operatorname{tr} A_i(b_n) \Delta_i s^2 + \operatorname{Scal}(\phi_n(s_{i-1})) \frac{1}{n} \right) = -2 \int_{s_{i-1}}^{s_i} \left\langle Ric_{u_n(s_{i-1})}(b(s) - b(s_{i-1})), db(s) \right\rangle.$$

Define

$$\begin{aligned} \xi_n &= 2 \sum_{i=1}^n \int_{s_{i-1}}^{s_i} \left\langle Ric_{u_n(s_{i-1})}(b(s) - b(s_{i-1})), db(s) \right\rangle \\ &= 2 \sum_{i=1}^n \left( \int_0^1 \left\langle 1_{J_i}(s) Ric_{u_n(s_{i-1})}(b(s) - b(s_{i-1})), db(s) \right\rangle \right) \\ &= 2 \int_0^1 \langle M_n(s), db(s) \rangle, \end{aligned}$$

where  $M_n(s) = \sum_{i=1}^n 1_{J_i}(s) Ric_{u_n(s_{i-1})}(b(s) - b(s_{i-1}))$ . To complete the proof, it suffices to show that  $\xi_n$  converges to 0  $\mu$ -a.s.. We will make use of Burkholder's Inequality,

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} |M_t|^p \right] \leq C \mathbb{E} \left[ \langle M \rangle_T^{\frac{p}{2}} \right],$$

where C is some constant. Thus applying this with  $T = 1$  and  $p = 4$ , we have

$$\begin{aligned} \mathbb{E} |\xi_n|^4 &\leq \mathbb{E} \left[ \left( C \int_0^1 \|M_n(s)\|^2 ds \right)^2 \right] \\ &\leq \mathbb{E} \left[ C^2 \int_0^1 \|M_n(s)\|^4 ds \right] \quad (\text{By Jensen's Inequality}) \\ &= \mathbb{E} \left[ C^2 \int_0^1 \sum_{i=1}^n 1_{J_i}(s) \|Ric_{u_n(s_{i-1})}(b(s) - b(s_{i-1}))\|^4 ds \right] \\ &= O \left( \int_0^1 \sum_{i=1}^n 1_{J_i}(s) \mathbb{E} \|b(s) - b(s_{i-1})\|^4 ds \right) \\ &= O \left( \int_0^1 \sum_{i=1}^n 1_{J_i}(s) (s - s_{i-1})^2 ds \right) = O \left( \frac{1}{n^2} \right). \end{aligned}$$

Hence  $\sum_{n=1}^{\infty} \mathbb{E} |\xi_n|^4 < \infty$  and thus

$$\xi_n \longrightarrow 0 \quad \mu - a.s..$$

■

**Proposition 5.10**

$$\sum_{i=1}^n Scal(\phi_n(s_{i-1})) \frac{1}{n} \longrightarrow \int_0^1 Scal(\tilde{\phi}(s)) ds$$

$\mu$ -a.s. as  $n \rightarrow \infty$ .

**Proof.** Note that we can write

$$\sum_{i=1}^n Scal(\phi_n(s_{i-1})) \frac{1}{n} = \int_0^1 \sum_{i=1}^n 1_{J_i}(s) Scal(\phi_n(s_{i-1})) ds.$$

Since  $\phi_n = \phi \circ b_n \rightarrow \tilde{\phi}$  in the sup norm  $\mu$ -a.s. as  $n \rightarrow \infty$  and  $Scal$  is a continuous function, thus

$$\begin{aligned} \sum_{i=1}^n 1_{J_i}(s) Scal(\phi_n(s_{i-1})) &= \sum_{i=1}^n 1_{J_i}(s) \left[ Scal(\phi_n(s_{i-1})) - Scal(\tilde{\phi}(s_{i-1})) + Scal(\tilde{\phi}(s_{i-1})) \right] \\ &\longrightarrow Scal(\tilde{\phi}(s)). \end{aligned}$$

Hence we can apply the dominated convergence theorem to obtain

$$\lim_{n \rightarrow \infty} \int_0^1 \sum_{i=1}^n 1_{J_i}(s) Scal(\phi_n(s_{i-1})) ds = \int_0^1 Scal(\tilde{\phi}(s)) ds$$

$\mu - a.s..$

■

**Lemma 5.11**

$$\det(\widetilde{\mathcal{V}^n \circ b_n}) = 1_{\{n \geq n_0\}} \prod_{i=1}^n \det \left[ I + \frac{1}{6} A_i(b_n) \Delta_i s^2 + \eta_i(b_n) \right] \longrightarrow e^{-\frac{1}{6} \int_0^1 \text{Scal}(\tilde{\phi}(s)) \, ds}$$

$\mu - a.s.$  as  $n \rightarrow \infty$ .

**Proof.** It suffices to consider on  $W_\alpha(\mathbb{R}^d)$  with  $\frac{1}{3} < \alpha < \frac{1}{2}$ , since  $\mu(W_\alpha(\mathbb{R}^d)) = 1$ . For  $n \geq n_0(\cdot)$ , write

$$\zeta_i^n = \frac{1}{6} A_i(b_n) \Delta_i s^2 + \eta_i(b_n)$$

where  $\eta_i(b_n)$  was defined in Equation (4.29) and  $\|\eta_i(b_n)\| = O\left(\bigvee_{i=1, \dots, n} \|\Delta_i b\|^3\right)$ . Now using the perturbation formula in Equation (B.2) of the Appendix with  $r = 2$ ,

$$\begin{aligned} \det[I + \zeta_i^n] &= \exp[\text{tr} \zeta_i^n + R_2(\zeta_i^n)] \\ &= \exp\left[\frac{1}{6} \text{tr} A_i(b_n) \Delta_i s^2 + \psi_i^n\right] \end{aligned}$$

where

$$\psi_i^n := \text{tr} \eta_i(b_n) + R_2(\zeta_i^n) = \text{tr} \eta_i(b_n) + \sum_{k=2}^{\infty} (-1)^{k+1} \text{tr} \left( \frac{1}{6} A_i(b_n) \Delta_i s^2 + \eta_i(b_n) \right)^k.$$

Using Equation (B.3),

$$|R_2(\zeta_i^n)| \leq \frac{d \|\zeta_i^n\|^2}{1 - \|\zeta_i^n\|} = O\left(\bigvee_{i=1, \dots, n} \|\Delta_i b\|^4\right)$$

and hence

$$|\psi_i^n| \leq |\text{tr} \eta_i(b_n)| + |R_2(\zeta_i^n)| = O\left(\bigvee_{i=1, \dots, n} \|\Delta_i b\|^3\right).$$

Since we chose  $\alpha > \frac{1}{3}$ , on  $W_\alpha(\mathbb{R}^d)$ ,

$$\begin{aligned} \left\| \sum_{i=1}^n \psi_i^n \right\| &= O\left(n \bigvee_{i=1, \dots, n} \|\Delta_i b\|^3\right) \\ &= O(n \cdot n^{-3\alpha}) = O(n^{1-3\alpha}) \\ &\longrightarrow 0 \end{aligned}$$

as  $n \rightarrow \infty$ . Together with Lemma 5.9,

$$\begin{aligned}
& 1_{\{n \geq n_0\}} \prod_{i=1}^n \det \left[ I + \frac{1}{6} A_i(b_n) \Delta_i s^2 + \eta_i(b_n) \right] - \exp \left[ -\frac{1}{6} \sum_{i=1}^n \text{Scal}(\phi_n(s_{i-1})) \frac{1}{n} \right] \\
&= 1_{\{n \geq n_0\}} \exp \left[ \frac{1}{6} \sum_{i=1}^n (\text{tr } A_i(b_n) \Delta_i s^2 + \psi_i^n) \right] - \exp \left[ -\frac{1}{6} \sum_{i=1}^n \text{Scal}(\phi_n(s_{i-1})) \frac{1}{n} \right] \\
&= e^{-\frac{1}{6} \sum_{i=1}^n \text{Scal}(\phi_n(s_{i-1})) \frac{1}{n}} \left\{ 1_{\{n \geq n_0\}} \exp \left[ \frac{1}{6} \sum_{i=1}^n \left( \text{tr } A_i(b_n) \Delta_i s^2 + \text{Scal}(\phi_n(s_{i-1})) \frac{1}{n} \right) + \sum_{i=1}^n \psi_i^n \right] - 1 \right\} \\
&\longrightarrow 0
\end{aligned}$$

$\mu$ -a.s. as  $n \rightarrow \infty$ . Finally,

$$\begin{aligned}
& 1_{\{n \geq n_0\}} \prod_{i=1}^n \det \left[ I + \frac{1}{6} A_i(b_n) \Delta_i s^2 + \eta_i(b_n) \right] - e^{-\frac{1}{6} \int_0^1 \text{Scal}(\tilde{\phi}(s)) ds} \\
&= 1_{\{n \geq n_0\}} \prod_{i=1}^n \det \left[ I + \frac{1}{6} A_i(b_n) \Delta_i s^2 + \eta_i(b_n) \right] - e^{-\frac{1}{6} \sum_{i=1}^n \text{Scal}(\phi_n(s_{i-1})) \frac{1}{n}} \\
&+ e^{-\frac{1}{6} \sum_{i=1}^n \text{Scal}(\phi_n(s_{i-1})) \frac{1}{n}} - e^{-\frac{1}{6} \int_0^1 \text{Scal}(\tilde{\phi}(s)) ds} \\
&\longrightarrow 0
\end{aligned}$$

$\mu$ -a.s. as  $n \rightarrow \infty$ . ■

## 5.2 Convergence of $\det(\mathcal{I}^n + \mathcal{U}^n) \circ b_n$

Recall, from Equation (4.36), that  $\mathcal{U}^n := (\mathcal{S}^n)^T \mathcal{C}^n \mathcal{S}^n$  where

$$\mathcal{C}_{ij}^n \circ b_n = \delta_{ij} \left( \frac{1}{45} (A_i^2(b_n) \Delta_i s^4 + A_{i+1}^2(b_n) \Delta_{i+1} s^4) \right) + 1_{\{|j-i|=1\}} \frac{7}{360} A_{i \vee j}^2(b_n) \Delta_{i \vee j} s^4$$

with  $A_{n+1}^2(b_n) \Delta_{n+1} s^4 := 0$  as in Equation (4.35). In order to compute the limit of  $\det(\mathcal{I}^n + \mathcal{U}^n)$  as  $n \rightarrow \infty$ , it will be necessary to compute

$$\text{Tr } ([\mathcal{U}^n]^k) = \text{Tr } [(\mathcal{S}^n)^T \mathcal{C}^n \mathcal{S}^n]^k = \text{Tr } [(\mathcal{S}^n)^T \mathcal{C}^n (\mathcal{B}^n \mathcal{C}^n)^{k-1} \mathcal{S}^n] = \text{Tr } ([\mathcal{B}^n \mathcal{C}^n]^k),$$

where  $\mathcal{B}^n := \mathcal{S}^n (\mathcal{S}^n)^T$ .

**Lemma 5.12** *The matrix,  $\mathcal{B}^n := \mathcal{S}^n (\mathcal{S}^n)^T$ , is given by*

$$\mathcal{B}_{lm}^n = (l \wedge m) I \text{ for } l, m = 1, 2, \dots, n. \tag{5.2}$$

Moreover,  $\mathcal{B}^n$  and  $\mathcal{S}^n$  satisfy the norm estimates,

$$\| \mathcal{S}^n \| = O(n) \text{ and } \| \mathcal{B}^n \| = O(n^2).$$

**Proof.** By definition,

$$\begin{aligned}\mathcal{B}_{lm}^n &= \sum_{k=1}^n \mathcal{S}_{lk}^n (\mathcal{S}_{mk}^n)^T = \sum_{k=1}^n 1_{l \geq k} 1_{m \geq k} I \\ &= \sum_{k=1}^n 1_{\{l \wedge m \geq k\}} I = (l \wedge m) I.\end{aligned}$$

Let  $\lambda_1 \geq \lambda_2 \dots \geq \lambda_{nd}$  be the eigenvalues of  $\mathcal{B}^n$ . Since it is a positive definite matrix, we have that  $\|\mathcal{B}^n\| = \lambda_1$ . Therefore, we have

$$\|\mathcal{B}^n\| = \lambda_1 \leq \sum_{i=1}^{nd} \lambda_i = \text{Tr } \mathcal{B}^n = \sum_{l=1}^n l \text{ tr } I = d \frac{n(n+1)}{2} = O(n^2).$$

Since

$$\begin{aligned}\|\mathcal{B}^n\| &= \sup_{\|v\|=1} \langle \mathcal{B}^n v, v \rangle = \sup_{\|v\|=1} \langle (\mathcal{S}^n)^T v, (\mathcal{S}^n)^T v \rangle \\ &= \|(\mathcal{S}^n)^T\|^2 = \|\mathcal{S}^n\|^2,\end{aligned}$$

it follows that  $\|\mathcal{S}^n\| = O(n)$ . ■

The following definition will be useful in describing the limiting behavior of  $A_m^2(b_n) \Delta_m s^4$  as  $n \rightarrow \infty$ .

**Definition 5.13** Define  $\Gamma : O(M) \rightarrow \mathbb{R}^{d \times d}$  (the  $d \times d$  matrices) by

$$\Gamma(v) = \sum_{i,j=1}^d \left( \Omega_v(e_i, \Omega_v(e_i, \cdot) e_j) e_j + \Omega_v(e_i, \Omega_v(e_j, \cdot) e_i) e_j + \Omega_v(e_i, \Omega_v(e_j, \cdot) e_j) e_i \right)$$

where  $\{e_i\}_{i=1,2,\dots,d}$  is any orthonormal basis for  $T_o M$ .

**Notation 5.14** For  $a_1, a_2, a_3, a_4 \in \mathbb{R}^d$  and  $1 \leq m \leq n$ , let  $\tilde{T}_m^n(a_1 \otimes a_2 \otimes a_3 \otimes a_4) : W(\mathbb{R}^d) \rightarrow \mathbb{R}^{d \times d}$  be defined by

$$\tilde{T}_m^n(a_1 \otimes a_2 \otimes a_3 \otimes a_4) := \Omega_{u_n(s_{m-1})}(a_1, \Omega_{u_n(s_{m-1})}(a_2, \cdot) a_3) a_4 \in \mathbb{R}^{d \times d}.$$

If  $\tau$  is a permutation of  $\{1, 2, 3, 4\}$ , let

$$(\tau \tilde{T}_m^n)(a_1 \otimes a_2 \otimes a_3 \otimes a_4) = \tilde{T}_m^n(a_{\tau(1)} \otimes a_{\tau(2)} \otimes a_{\tau(3)} \otimes a_{\tau(4)})$$

and

$$(\tau \tilde{T}_{m,i}^n)(a_1, a_2) = (\tau \tilde{T}_m^n)(e_i \otimes e_i \otimes a_1 \otimes a_2)$$

where  $\{e_i\}_{i=1}^d$  is the standard orthonormal basis for  $\mathbb{R}^d$ .

Let  $\underline{0} = 0$  and for  $t \in (s_{m-1}, s_m]$ , let  $\underline{t} = s_{m-1}$ ,  $\Delta := s_m - s_{m-1} = 1/n$ ,

$$\Delta b(t) = b(t) - b(\underline{t}),$$

$$\Delta_m b = \Delta b(s_m) = b(s_m) - b(s_{m-1}),$$

$$\Delta_m b^i = \Delta_m b \cdot e_i, \text{ and}$$

$$\Delta b(t)^{\otimes 3} \otimes db(t) = \Delta b(t) \otimes \Delta b(t) \otimes \Delta b(t) \otimes db(t).$$

**Lemma 5.15** *Let  $\{e_i\}_{i=1,2,\dots,d}$  be an orthonormal basis for  $T_o M$ ,  $\Delta = \frac{1}{n}$  and  $m \in \{1, 2, \dots, n\}$ . Then*

$$A_m^2(b_n)\Delta^4 - \Delta^2 \Gamma(u_n(s_{m-1})) = \epsilon_m^n \quad (5.3)$$

where

$$\epsilon_m^n := \frac{1}{6} \int_{s_{m-1}}^{s_m} \sum_{\tau} (\tau \tilde{T}_m^n)(\Delta b(s)^{\otimes 3} \otimes db(s)) + \frac{1}{2} \int_{s_{m-1}}^{s_m} (s_m - s) \sum_{i,\tau} (\tau \tilde{T}_{m,i}^n)(\Delta b(s) \otimes db(s)).$$

The above sums range over permutations,  $\tau$ , of  $\{1, 2, 3, 4\}$  and  $i = 1, 2, \dots, d$ .

**Proof.** By definition, we have

$$\begin{aligned} A_m^2(b_n)\Delta^4 &= \Omega_{u_n(s_{m-1})}(\Delta_m b, \Omega_{u_n(s_{m-1})}(\Delta_m b, \cdot) \Delta_m b) \Delta_m b \\ &= \tilde{T}_m^n(\Delta_m b \otimes \Delta_m b \otimes \Delta_m b \otimes \Delta_m b) \\ &= \sum_{i,j,k,l} \tilde{T}_m^n(e_i \otimes e_j \otimes e_k \otimes e_l) \Delta_m b^i \Delta_m b^j \Delta_m b^k \Delta_m b^l. \end{aligned}$$

By Ito's formula,

$$\begin{aligned} &\tilde{T}_m^n(\Delta_m b \otimes \Delta_m b \otimes \Delta_m b \otimes \Delta_m b) \\ &= \int_{s_{m-1}}^{s_m} \sum_{j=1}^4 \tilde{T}_m^n(\Delta b(s) \otimes \dots \underbrace{db(s)}_{j^{\text{th}} \text{ - spot}} \dots \otimes \Delta b(s)) + \frac{1}{4} \int_{s_{m-1}}^{s_m} \sum_{i=1}^d \sum_{\tau} (\tau \tilde{T}_m^n)(e_i \otimes e_i \otimes \Delta b(s) \otimes \Delta b(s)) ds \\ &= \frac{1}{6} \int_{s_{m-1}}^{s_m} \sum_{\tau} (\tau \tilde{T}_m^n)(\Delta b(s)^{\otimes 3} \otimes db(s)) + \frac{1}{4} \int_{s_{m-1}}^{s_m} \sum_{i,\tau} (\tau \tilde{T}_{m,i}^n)(\Delta b(s) \otimes \Delta b(s)) ds. \end{aligned} \quad (5.4)$$

Now we claim that

$$\int_{s_{m-1}}^{s_m} \Delta b(s) \otimes \Delta b(s) ds = \int_{s_{m-1}}^{s_m} (s_m - t) \Delta b(t) \vee db(t) + \frac{\Delta^2}{2} \sum_{i=1}^d e_i \otimes e_i, \quad (5.5)$$

where  $a_1 \vee a_2 = a_1 \otimes a_2 + a_2 \otimes a_1$ . Let

$$\begin{aligned} W_s &= \int_{s_{m-1}}^s (s - t) \Delta b(t) \vee db(t) + \frac{(s - s_{m-1})^2}{2} \sum_{i=1}^d e_i \otimes e_i \\ &= s \int_{s_{m-1}}^s \Delta b(t) \vee db(t) - \int_{s_{m-1}}^s t \Delta b(t) \vee db(t) + \frac{(s - s_{m-1})^2}{2} \sum_{i=1}^d e_i \otimes e_i \end{aligned} \quad (5.6)$$

and observe that  $W_{s_m}$  is equal to the right side of Equation (5.5). Since,

$$\begin{aligned} dW_s &= \left( \int_{s_{m-1}}^s \Delta b(t) \vee db(t) \right) ds + s \Delta b(s) \vee db(s) - s \Delta b(s) \vee db(s) + \left( (s - s_{m-1}) \sum_{i=1} e_i \otimes e_i \right) ds \\ &= \left( \int_{s_{m-1}}^s \Delta b(t) \vee db(t) + (s - s_{m-1}) \sum_{i=1} e_i \otimes e_i \right) ds \\ &= (\Delta b(s) \otimes \Delta b(s)) ds, \end{aligned}$$

it follows that

$$W_{s_m} = \int_{s_{m-1}}^{s_m} \Delta b(s) \otimes \Delta b(s) ds$$

which verifies Equation (5.5).

From Equation (5.5), we have

$$\begin{aligned} & \int_{s_{m-1}}^{s_m} \sum_{i,\tau} (\tau \tilde{T}_{m,i}^n) (\Delta b(s) \otimes \Delta b(s)) ds \\ &= \int_{s_{m-1}}^{s_m} (s_m - s) \sum_{i,\tau} (\tau \tilde{T}_{m,i}^n) (\Delta b(s) \vee db(s)) + \frac{\Delta^2}{2} \sum_{i,\tau} \sum_{j=1}^d (\tau \tilde{T}_{m,i}^n) (e_j \otimes e_j) \\ &= \int_{s_{m-1}}^{s_m} (s_m - s) \sum_{i,\tau} (\tau \tilde{T}_{m,i}^n) (\Delta b(s) \vee db(s)) + \frac{\Delta^2}{2} \sum_{\tau} \sum_{i,j=1}^d (\tau \tilde{T}_m^n) (e_i \otimes e_i \otimes e_j \otimes e_j) \\ &= 2 \int_{s_{m-1}}^{s_m} (s_m - s) \sum_{i,\tau} (\tau \tilde{T}_{m,i}^n) (\Delta b(s) \otimes db(s)) + \frac{\Delta^2}{2} 8 \Gamma(u(s_{m-1})) \\ &= 2 \int_{s_{m-1}}^{s_m} (s_m - s) \sum_{i,\tau} (\tau \tilde{T}_{m,i}^n) (\Delta b(s) \otimes db(s)) + 4 \Delta^2 \Gamma(u_n(s_{m-1})). \end{aligned}$$

When we sum over all permutations of  $\{1, 2, 3, 4\}$ , we will have  $4! = 24$  terms. However,  $\Gamma$  is a sum of 3 terms, hence we will end up with 8 copies of  $\Gamma$ , which explains the factor 8 in the second last equality. Combining this equation with Equation (5.4) proves Equation (5.3).  $\blacksquare$

**Lemma 5.16** *There exists  $C < \infty$  such that for any  $\lambda > 0$ ,  $n \in \mathbb{N}$ , and  $\{c_m\}_{m=1}^n \subset [0, \lambda]$ ,*

$$\mathbb{E} \left\| \sum_{m=1}^n c_m \epsilon_m^n \right\|^2 \leq C \frac{\lambda^2}{n^3},$$

which as usual we abbreviate as

$$\mathbb{E} \left\| \sum_{m=1}^n c_m \epsilon_m^n \right\|^2 = O\left(\frac{\lambda^2}{n^3}\right).$$

**Proof.** We may write

$$\sum_{m=1}^n c_m \epsilon_m^n = \xi_n + \chi_n$$



where

$$\xi_n = \frac{1}{6} \sum_{m=1}^n c_m \int_{s_{m-1}}^{s_m} \sum_{\tau} (\tau \tilde{T}_m^n)(\Delta b(s))^{\otimes 3} \otimes db(s)$$

and

$$\chi_n = \frac{1}{2} \sum_{m=1}^n c_m \int_{s_{m-1}}^{s_m} (s_m - s) \sum_{i, \tau} (\tau \tilde{T}_{m,i}^n)(\Delta b(s) \otimes db(s)).$$

Using the isometry property of the Ito integral, we have

$$\begin{aligned} \mathbb{E} (6\xi_n)^2 &= \mathbb{E} \left\| \sum_{m=1}^n c_m \int_{s_{m-1}}^{s_m} \sum_{\tau} (\tau \tilde{T}_m^n)(\Delta b(s))^{\otimes 3} \otimes db(s) \right\|^2 \\ &= \mathbb{E} \left\| \int_0^1 \sum_{m=1}^n c_m 1_{J_m}(s) \sum_{\tau} (\tau \tilde{T}_m^n)(\Delta b(s))^{\otimes 3} \otimes db(s) \right\|^2 \\ &= O \left( \mathbb{E} \int_0^1 \sum_{m=1}^n c_m^2 1_{J_m}(s) \|\Delta b(s)\|^6 ds \right) \\ &= O \left( \int_0^1 \sum_{m=1}^n c_m^2 1_{J_m}(s) (s - s_{m-1})^3 ds \right) \\ &= O \left( \sum_{m=1}^n c_m^2 \frac{1}{n^4} \right) = O \left( \frac{\lambda^2}{n^3} \right). \end{aligned}$$

Similarly,

$$\begin{aligned} \mathbb{E} (2\chi_n)^2 &= \mathbb{E} \left\| \sum_{m=1}^n c_m \int_{s_{m-1}}^{s_m} (s_m - s) \sum_{i, \tau} (\tau \tilde{T}_{m,i}^n)(\Delta b(s) \otimes db(s)) \right\|^2 \\ &= \mathbb{E} \left\| \int_0^1 \sum_{m=1}^n c_m 1_{J_m}(s) (s_m - s) \sum_{i=1}^d \sum_{\tau} (\tau \tilde{T}_{m,i}^n)(\Delta b(s) \otimes db(s)) \right\|^2 \\ &= O \left( \mathbb{E} \int_0^1 \sum_{m=1}^n c_m^2 1_{J_m}(s) (s_m - s)^2 \|\Delta b(s)\|^2 ds \right) \\ &= O \left( \int_0^1 \sum_{m=1}^n c_m^2 1_{J_m}(s) (s_m - s)^2 (s - s_{m-1}) ds \right) \\ &= O \left( \sum_{m=1}^n c_m^2 \frac{1}{n^4} \right) = O \left( \frac{\lambda^2}{n^3} \right). \end{aligned}$$

Thus

$$\left( \mathbb{E} \left\| \sum_{m=1}^n c_m \epsilon_m^n \right\|^2 \right)^{\frac{1}{2}} \leq \left( \mathbb{E} \|\xi_n\|^2 \right)^{\frac{1}{2}} + \left( \mathbb{E} \|\chi_n\|^2 \right)^{\frac{1}{2}} = O \left( \sqrt{\frac{\lambda^2}{n^3}} \right)$$

and hence

$$\mathbb{E} \left\| \sum_{m=1}^n c_m \epsilon_m^n \right\|^2 = O \left( \frac{\lambda^2}{n^3} \right).$$

■

**Definition 5.17** For  $l, m = 1, 2, \dots, n$ , define  $\Lambda_{lm}^n$  and  $\Phi_{lm}^n$  by

$$\begin{aligned}\Lambda_{lm}^n &= \left(l \wedge (m-1)\right) \frac{7}{360} \Gamma(u_n(s_{m-1})) \frac{1}{n^2} + \left(l \wedge m\right) \frac{1}{45} \Gamma(u_n(s_m)) \frac{1}{n^2} 1_{\{m < n\}} \\ &\quad + \left(l \wedge m\right) \frac{1}{45} \Gamma(u_n(s_{m-1})) \frac{1}{n^2} + \left(l \wedge (m+1)\right) \frac{7}{360} \Gamma(u_n(s_m)) \frac{1}{n^2} 1_{\{m < n\}}\end{aligned}$$

and

$$\begin{aligned}\Phi_{lm}^n &= \left(l \wedge (m-1)\right) \frac{7}{360} \epsilon_m^n + \left(l \wedge m\right) \frac{1}{45} \epsilon_{m+1}^n 1_{\{m < n\}} \\ &\quad + \left(l \wedge m\right) \frac{1}{45} \epsilon_m^n + \left(l \wedge (m+1)\right) \frac{7}{360} \epsilon_{m+1}^n 1_{\{m < n\}}.\end{aligned}$$

With this notation along with Equation (5.2) and Equation (5.3), we have

$$\begin{aligned}(\mathcal{B}^n \mathcal{C}^n)_{lm} \circ b_n &= \left( \sum_{k=1}^n \mathcal{B}_{lk}^n \mathcal{C}_{km}^n \right) \circ b_n = \left( \sum_{k=1}^n (l \wedge k) \mathcal{C}_{km}^n \right) \circ b_n \\ &= \Lambda_{lm}^n + \Phi_{lm}^n.\end{aligned}\tag{5.7}$$

**Notation 5.18** For any  $k \in \mathbb{N}$  and  $d \times d$  matrices  $\{M_k\}_{k=1}^k$ , let

$$\left[ \prod_{l=1}^k \right] M_k := M_1 M_2 \cdots M_k.$$

**Theorem 5.19** For each  $k = 1, 2, \dots$ , define  $\gamma_k^n : W(\mathbb{R}^d) \rightarrow \mathbb{R}$  by

$$\gamma_k^n := \sum_{r_k=1}^n \cdots \sum_{r_1=1}^n \text{tr} \left[ \prod_{l=1}^k \right] \Lambda_{r_i, r_{i+1}}^n,$$

where  $r_{k+1} = r_1$ . Then

$$\lim_{n \rightarrow \infty} \mathbb{E} \left| \text{Tr} ([\mathcal{U}^n \circ b_n]^k) - \gamma_k^n \right| = 0$$

and in particular

$$\text{Tr} ([\mathcal{U}^n \circ b_n]^k) - \gamma_k^n \longrightarrow 0 \text{ (in } \mu \text{ - measure) as } n \rightarrow \infty.$$

**Proof.** For a fixed  $k \in \mathbb{N}$ ,

$$\begin{aligned}\text{Tr} ([\mathcal{B}^n \mathcal{C}^n \circ b_n]^k) &= \left( \text{tr} \sum_{r_k=1}^n \cdots \sum_{r_1=1}^n (\mathcal{B}^n \mathcal{C}^n)_{r_1, r_2} (\mathcal{B}^n \mathcal{C}^n)_{r_2, r_3} \cdots (\mathcal{B}^n \mathcal{C}^n)_{r_k, r_1} \right) \circ b_n \\ &= \text{tr} \sum_{r_k=1}^n \cdots \sum_{r_1=1}^n \Lambda_{r_1, r_2}^n \Lambda_{r_2, r_3}^n \cdots \Lambda_{r_k, r_1}^n + \Xi^n \\ &= \gamma_k^n + \Xi^n,\end{aligned}$$

where  $\Xi^n$  consists of a finite sum of terms of the form

$$\text{tr} \sum_{r_k=1}^n \cdots \sum_{r_1=1}^n c_{r_1, r_2} c_{r_2, r_3} \cdots c_{r_k, r_1} \theta_{r_1} \theta_{r_2} \cdots \theta_{r_k},$$

with

$$\theta_{r_i} \in \left\{ \frac{1}{n^2} \Gamma(u_n(s_{r_i-1})), \frac{1}{n^2} \Gamma(u_n(s_{r_i})), \epsilon_{r_i}^n, \epsilon_{r_i+1}^n \right\},$$

$1 \leq c_{r_i, r_{i+1}} \leq n$  for  $i \leq n$  and for at least one  $r_i$ ,  $\theta_{r_i} = \epsilon_{r_i}^n$ . Since trace is invariant under cyclic permutation, we can assume that  $\theta_{r_k} = \epsilon_{r_k}^n$ . To finish the proof it suffices to show,  $\lim_{n \rightarrow \infty} \mathbb{E} \|\Xi^n\| = 0$  and for this it suffices to show,

$$\lim_{n \rightarrow \infty} \mathbb{E} \left\| \sum_{r_k=1}^n \cdots \sum_{r_1=1}^n c_{r_1, r_2} c_{r_2, r_3} \cdots c_{r_k, r_1} \theta_{r_1} \theta_{r_2} \cdots \theta_{r_k} \right\| = 0.$$

Let  $r = (r_1, \dots, r_k)$  and write

$$c(r) = \prod_{i=1}^k c_{r_i, r_{i+1}},$$

where by convention,  $r_{k+1} := r_1$ . Then

$$\begin{aligned} T_n &:= \left( \mathbb{E} \left\| \sum_{r_k=1}^n \cdots \sum_{r_1=1}^n c(r) \theta_{r_1} \theta_{r_2} \cdots \theta_{r_{k-1}} \epsilon_{r_k} \right\| \right)^2 \\ &= \left( \mathbb{E} \left\| \sum_{r_{k-1}=1}^n \cdots \sum_{r_1=1}^n \theta_{r_1} \theta_{r_2} \cdots \theta_{r_{k-1}} \sum_{r_k=1}^n c(r) \epsilon_{r_k} \right\| \right)^2 \\ &\leq \left( \mathbb{E} \sum_{r_{k-1}=1}^n \cdots \sum_{r_1=1}^n \left( \|\theta_{r_1} \theta_{r_2} \cdots \theta_{r_{k-1}}\| \left\| \sum_{r_k=1}^n c(r) \epsilon_{r_k} \right\| \right) \right)^2 \\ &\leq \mathbb{E} \left( \sum_{r_1, \dots, r_{k-1}} \prod_{i=1}^{k-1} \|\theta_{r_i}\|^2 \right) \mathbb{E} \left( \sum_{r_1, \dots, r_{k-1}} \left\| \sum_{r_k=1}^n c(r) \epsilon_{r_k} \right\|^2 \right), \end{aligned} \quad (5.8)$$

where the last inequality is a consequence of the Cauchy Schwartz inequality. Let  $C_1 := \sup_{v \in O(M)} \|\Gamma(v)\| < \infty$ , then

$$\left\| \frac{1}{n^2} \Gamma(u(s_{i-1})) \right\|^{2(k-1)} \leq C_1^{2(k-1)} \left( \frac{1}{n} \right)^{4(k-1)} = O\left( \frac{1}{n^{4(k-1)}} \right).$$

Using Equation (5.3), we observe that there exists a constant  $C_2$  such that

$$\mathbb{E} \|\epsilon_m^n\|^{2(k-1)} \leq \mathbb{E} \left[ C_2 \left( \|\Delta_m b\|^4 + \frac{1}{n^2} \right) \right]^{2(k-1)} = O\left( \frac{1}{n^{4(k-1)}} \right).$$

Thus we can find a constant  $C(k)$  such that

$$\mathbb{E} \|\theta_{r_i}\|^{2(k-1)} \leq C \left( \frac{1}{n} \right)^{4(k-1)}.$$

By Holder's Inequality,

$$\begin{aligned} \mathbb{E} \left( \sum_{r_{k-1}=1}^n \cdots \sum_{r_1=1}^n \prod_{i=1}^{k-1} \|\theta_{r_i}\|^2 \right) &\leq \sum_{r_{k-1}=1}^n \cdots \sum_{r_1=1}^n \prod_{i=1}^{k-1} \left( \mathbb{E} \|\theta_{r_i}\|^{2(k-1)} \right)^{\frac{1}{k-1}} \\ &= \sum_{r_{k-1}=1}^n \cdots \sum_{r_1=1}^n C \left( \frac{1}{n} \right)^{4(k-1)} = C \left( \frac{1}{n} \right)^{3(k-1)}. \end{aligned} \quad (5.9)$$

Note that  $\sup_r |c(r)| \leq n^k$ . Using Lemma 5.16, with  $\lambda = n^k$ , we have

$$\begin{aligned} \mathbb{E} \left( \sum_{r_{k-1}=1}^n \cdots \sum_{r_1=1}^n \left\| \sum_{r_k=1}^n c(r) \epsilon_{r_k} \right\|^2 \right) &= \sum_{r_{k-1}=1}^n \cdots \sum_{r_1=1}^n \mathbb{E} \left\| \sum_{r_k=1}^n c(r) \epsilon_{r_k} \right\|^2 \\ &= \sum_{r_{k-1}=1}^n \cdots \sum_{r_1=1}^n O \left( \frac{n^{2k}}{n^3} \right) = n^{k-1} O \left( \frac{n^{2k}}{n^3} \right) = n^{3(k-1)} O \left( \frac{1}{n} \right). \end{aligned} \quad (5.10)$$

Hence using Equations (5.9) and (5.10), we have from Equation (5.8),

$$T_n \leq C \frac{1}{n^{3(k-1)}} n^{3(k-1)} \frac{1}{n} = O \left( \frac{1}{n} \right).$$

Thus putting all together, we have

$$\mathbb{E} \|\Xi^n\| \leq d \cdot \sqrt{T_n} = O \left( \frac{1}{\sqrt{n}} \right).$$

■

**Definition 5.20** *Let*

$$h(s, t) = (s \wedge t) \Gamma(u(t))$$

*and for  $k \in \mathbb{N}$ , let*

$$\gamma_k = \left( \frac{1}{12} \right)^k \int_{s_k=0}^1 \cdots \int_{s_1=0}^1 \text{tr} \left[ \prod_{l=1}^k \right] h(s_l, s_{l+1}) ds_1 \dots ds_k,$$

*where again by convention,  $s_{k+1} := s_1$ .*

**Proposition 5.21** *Continuing the notation in the above definition,*

$$\gamma_k^n \longrightarrow \gamma_k$$

*$\mu$ -a.s. as  $n \rightarrow \infty$ .*

**Proof.** To begin with, let us write

$$s = (s_1, s_2, \dots, s_k) \in [0, 1]^k,$$

$$r = (r_1, r_2, \dots, r_k) \in W_n^k := \{1, 2, \dots, n\}^k,$$

$$ds = ds_1 \dots ds_k$$

and

$$H(s) = \left[ \prod_{l=1}^k \right] h(s_l, s_{l+1})$$

where  $s_{k+1} = s_1$ . Denote  $r_l^- := r_l - 1$  and

$$V_n(r) = V_n(r_1, r_2, \dots, r_k) := J_{r_1} \times J_{r_2} \cdots J_{r_k} = \left( \frac{r_1^-}{n}, \frac{r_1}{n} \right] \times \left( \frac{r_2^-}{n}, \frac{r_2}{n} \right] \cdots \left( \frac{r_k^-}{n}, \frac{r_k}{n} \right].$$

With this new notation and  $r_{k+1} = r_1$ ,

$$\begin{aligned} \gamma_k^n - \gamma_k &= \sum_{r \in W_n^k} \text{tr} \left[ \prod_{l=1}^k \right] \Lambda_{r_i, r_{i+1}}^n - \left( \frac{1}{12} \right)^k \int_{s \in [0,1]^k} \text{tr} H(s) ds \\ &= \sum_{r \in W_n^k} \left( \text{tr} \left[ \prod_{l=1}^k \right] \Lambda_{r_i, r_{i+1}}^n - \left( \frac{1}{12} \right)^k \int_{s \in V_n(r)} \text{tr} H(s) ds \right) \\ &= \sum_{r \in W_n^k} \int_{s \in V_n(r)} \left( n^k \text{tr} \left[ \prod_{l=1}^k \right] \Lambda_{r_i, r_{i+1}}^n - \left( \frac{1}{12} \right)^k \text{tr} H(s) \right) ds \\ &= \int_{s \in [0,1]^k} \sum_{r \in W_n^k} 1_{V_n(r)}(s) \left( \text{tr} \left[ \prod_{l=1}^k \right] n \Lambda_{r_i, r_{i+1}}^n - \left( \frac{1}{12} \right)^k \text{tr} H(s) \right) ds. \end{aligned}$$

Now  $u_n \rightarrow u$  in the sup norm  $\mu$ -a.s. and  $\Gamma(\cdot)$  is continuous, thus

$$\begin{aligned} \sum_{l,m=1}^n 1_{J_l \times J_m}(\tau, t) \left( \frac{l}{n} \wedge \frac{m-1}{n} \right) \Gamma \left( u_n \left( \frac{m-1}{n} \right) \right) - h(\tau, t) &\longrightarrow 0 \text{ and} \\ \sum_{l,m=1}^n 1_{J_l \times J_m}(\tau, t) \left( \frac{l}{n} \wedge \frac{m}{n} \right) \Gamma \left( u_n \left( \frac{m}{n} \right) \right) - h(\tau, t) &\longrightarrow 0 \end{aligned}$$

$\mu$ -a.s. as  $n \rightarrow \infty$ . For  $(\tau, t) \in [0, 1]^2$ , by using Definition 5.17,

$$\begin{aligned} &\sum_{l,m=1}^n 1_{J_l \times J_m}(\tau, t) n \Lambda_{l,m}^n - \frac{1}{12} h(\tau, t) \\ &= \sum_{l,m=1}^n 1_{J_l \times J_m}(\tau, t) \left\{ \frac{7}{360} \left( \frac{l}{n} \wedge \frac{m-1}{n} \right) \Gamma \left( u_n \left( \frac{m-1}{n} \right) \right) + \frac{1}{45} \left( \frac{l}{n} \wedge \frac{m}{n} \right) \Gamma \left( u_n \left( \frac{m}{n} \right) \right) 1_{\{m < n\}} \right\} \\ &\quad - \frac{1}{12} h(\tau, t) \\ &\longrightarrow 0 \end{aligned}$$

$\mu$ -a.s. as  $n \rightarrow \infty$ . And since taking trace and products are continuous operations, for  $s \in [0, 1]^k$ , we hence have

$$\begin{aligned} g_n(s) &:= \sum_{r \in W_n^k} 1_{V_n(r)}(s) \left( \text{tr} \left[ \prod_{l=1}^k \right] n \Lambda_{r_i, r_{i+1}}^n - \left( \frac{1}{12} \right)^k \text{tr} H(s) \right) \\ &= \sum_{r \in W_n^k} 1_{V_n(r)}(s) \left( \text{tr} \left[ \prod_{l=1}^k \right] n \Lambda_{r_i, r_{i+1}}^n - \text{tr} \left[ \prod_{l=1}^k \right] \frac{1}{12} h(s_l, s_{l+1}) \right) \longrightarrow 0 \end{aligned}$$

$\mu$ -a.s. as  $n \rightarrow \infty$ . Thus we can now apply dominated convergence theorem and this gives us

$$\gamma_k^n - \gamma_k = \int_{s \in [0,1)^k} g_n(s) ds \longrightarrow 0$$

as  $n \rightarrow \infty$ . ■

**Remark 5.22** *If  $S_v$  is the sectional curvature of the manifold and  $\sup_{v \in O(M)} \|S_v\| < \frac{3}{17d}$ , then for any orthonormal frame  $\{e_i\}_{i=1}^d \subseteq T_o M$ ,*

$$\sup_{v \in O(M)} \|\Omega_v(e_i, \cdot)e_j\| < 2/d,$$

and hence

$$\sup_{v \in O(M)} \|\Gamma(v)\| < 3d^2 \cdot 4/d^2 = 12.$$

See Definition D.1 and Proposition D.2. Let  $\kappa = \sup_{v \in O(M)} \|\Gamma(v)\| / 12$ . Then  $\kappa < 1$  and hence

$$|\gamma_k^n| \leq \sum_{r_k=1}^n \cdots \sum_{r_1=1}^n d \left( \frac{1}{12n} \right)^k \sup_{v \in O(M)} \|\Gamma(v)\|^k = d\kappa^k.$$

A standard result in probability states that  $x_n \rightarrow x$  in  $\mu$ -measure iff for every subsequence of  $\{x_n, n \geq 1\}$  has itself a subsequence converging  $\mu$ -a.s. to  $x$ . Hence for any continuous function  $f$ ,  $f(x_n)$  converges to  $f(x)$  in  $\mu$ -measure. In the next proof, we will be using this result without further comment.

**Theorem 5.23** *If  $\sup_{v \in O(M)} \|S_v\| < \frac{3}{17d}$ , then*

$$\det[(\mathcal{I}^n + \mathcal{U}^n) \circ b_n] \longrightarrow e^\gamma$$

in  $\mu$ -measure as  $n \rightarrow \infty$ , where  $\gamma$  is defined as

$$\gamma := \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k} \gamma_k.$$

**Proof.** To ease the notation, in this proof only, we will write  $\mathcal{U}^n = \mathcal{U}^n \circ b_n$  and  $\mathcal{I}^n = \mathcal{I}^n \circ b_n$ . By Remark 5.22,  $|\gamma_k^n| \leq d\kappa^k$ ,  $\kappa = \sup_{v \in O(M)} \|\Gamma(v)\| / 12 < 1$ . Observe that  $\mathcal{U}^n$  satisfies the hypothesis in Lemma B.1. Thus we can apply the formula in Equation (B.1). Hence

$$\det(\mathcal{I}^n + \mathcal{U}^n) = \exp(\Psi_r + R_{r+1}),$$

where  $\Psi_r = \sum_{k=1}^r (-1)^{k+1} \frac{1}{k} \text{Tr} ([\mathcal{U}^n]^k)$  and  $|R_{r+1}| \leq \frac{1}{r+1} \text{Tr} ([\mathcal{U}^n]^{r+1})$ . Therefore

$$\begin{aligned} & \left| \det(\mathcal{I}^n + \mathcal{U}^n) - \exp \left( \sum_{k=1}^r (-1)^{k+1} \frac{1}{k} \gamma_k^n \right) \right| \\ &= \exp \left( \sum_{k=1}^r \frac{(-1)^{k+1}}{k} \gamma_k^n \right) \left| \exp \left( \sum_{k=1}^r \frac{(-1)^{k+1}}{k} (\text{Tr} ([\mathcal{U}^n]^k) - \gamma_k^n) + R_{r+1} \right) - 1 \right| \\ &\leq \exp(d\kappa/(1-\kappa)) \left| \exp \left( \sum_{k=1}^r (-1)^{k+1} \frac{1}{k} (\text{Tr} ([\mathcal{U}^n]^k) - \gamma_k^n) + R_{r+1} \right) - 1 \right|. \end{aligned}$$

The last inequality follows from

$$\exp \left( \sum_{k=1}^r (-1)^{k+1} \frac{1}{k} \gamma_k^n \right) \leq \exp \left( \sum_{k=1}^r \frac{1}{k} d\kappa^k \right) < \exp(d\kappa/(1-\kappa)).$$

Now by Theorem 5.19, we have

$$\text{Tr} ([\mathcal{U}^n]^k) - \gamma_k^n \rightarrow 0$$

in  $\mu$ -measure as  $n \rightarrow \infty$ . Together with  $\|\gamma_k^n\| \leq d\kappa^k$ , we will have

$$\limsup_{n \rightarrow \infty} |R_{r+1}| \leq \frac{1}{r+1} \limsup_{n \rightarrow \infty} \text{Tr} ([\mathcal{U}^n]^{r+1}) \leq \frac{1}{(r+1)} d\kappa^{r+1} \leq d\kappa^{r+1},$$

in  $\mu$ -measure. Therefore

$$\limsup_{n \rightarrow \infty} \left| \det[\mathcal{I}^n + \mathcal{U}^n] - \exp \left( \sum_{k=1}^r (-1)^{k+1} \frac{1}{k} \gamma_k^n \right) \right| \leq \exp(d\kappa/(1-\kappa)) |\exp(d\kappa^{r+1}) - 1| \quad (5.11)$$

in  $\mu$ -measure. From Proposition 5.21, we know that  $\gamma_k^n \rightarrow \gamma_k$  and hence  $\gamma_k \leq d\kappa^k$ , which implies

$p_r = \sum_{k=1}^r \frac{(-1)^{k+1}}{k} \gamma_k$  converges to  $\gamma$  and  $|p_r| \leq \frac{d\kappa}{1-\kappa}$ . Thus

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left| e^\gamma - \exp \left( \sum_{k=1}^r \frac{(-1)^{k+1}}{k} \gamma_k^n \right) \right| \leq |e^\gamma - e^{p_r}| + \lim_{n \rightarrow \infty} \left| e^{p_r} - \exp \left( \sum_{k=1}^r \frac{(-1)^{k+1}}{k} \gamma_k^n \right) \right| \\ &\leq e^{p_r} \left| \exp \left( \sum_{k=r+1}^{\infty} (-1)^{k+1} \frac{1}{k} \gamma_k \right) - 1 \right| \leq e^{p_r} \left| \exp \left( \sum_{k=r+1}^{\infty} \frac{1}{k} d\kappa^k \right) - 1 \right| \\ &\leq e^{d\kappa/(1-\kappa)} \left| \exp \left( \frac{d\kappa^{r+1}}{1-\kappa} \right) - 1 \right| \end{aligned} \quad (5.12)$$

$\mu$ -a.s.. Therefore using Equations (5.11) and (5.12),

$$\begin{aligned} & \limsup_{n \rightarrow \infty} |\det[\mathcal{I}^n + \mathcal{U}^n] - e^\gamma| \\ &\leq \limsup_{n \rightarrow \infty} \left\{ \left| e^\gamma - \exp \left( \sum_{k=1}^r \frac{(-1)^{k+1}}{k} \gamma_k^n \right) \right| + \left| \det[\mathcal{I}^n + \mathcal{U}^n] - \exp \left( \sum_{k=1}^r \frac{(-1)^{k+1}}{k} \gamma_k^n \right) \right| \right\} \\ &\leq \exp(d\kappa/(1-\kappa)) \left| \exp \left( \frac{d\kappa^{r+1}}{1-\kappa} \right) - 1 \right| + \exp(d\kappa/(1-\kappa)) |\exp(d\kappa^{r+1}) - 1| \end{aligned}$$

in  $\mu$ -measure. Since the inequality holds for every  $r$  and as  $r \rightarrow \infty$ ,

$$|\exp(a\kappa^{r+1}) - 1| \rightarrow 0,$$

for any constant  $a$ , we thus have

$$\limsup_{n \rightarrow \infty} |\det[(\mathcal{I}^n + \mathcal{U}^n) \circ b_n] - e^\gamma| = 0$$

in  $\mu$ -measure. ■

**Definition 5.24** Define an integral operator  $K_u : L^2([0, 1] \rightarrow \mathbb{R}^d) \longrightarrow L^2([0, 1] \rightarrow \mathbb{R}^d)$  by

$$\begin{aligned} (K_u f)(s) &:= \int_0^1 h(s, t) f(t) dt \\ &= \int_0^1 (s \wedge t) \Gamma(u(t)) f(t) dt, \end{aligned}$$

where  $\Gamma$  was defined in Definition 5.13.

**Proposition 5.25**  $K_u$  is trace class.

**Proof.** Note that  $L^2([0, 1] \rightarrow \mathbb{R}^d)$  is a separable Hilbert space. Let  $(Af)(s) = \int_0^1 (s \wedge t) f(t) dt$  and  $(Bf)(s) = \Gamma(u(s)) f(s)$ . Thus  $K_u = AB$ . By Proposition A.2,  $A$  is trace class. Since  $B$  is bounded, by Proposition A.3,  $\text{tr } |AB| < \infty$  and hence  $K_u$  is trace class. ■

**Proposition 5.26**

$$e^\gamma = \det \left( I + \frac{1}{12} K_u \right).$$

**Proof.** We will use Equation (B.2) to prove the statement. Note that by Remark 5.22, for any  $f \in L^2([0, 1] \rightarrow \mathbb{R}^d)$ ,

$$\begin{aligned} \left\| \int_0^1 \frac{1}{12} (K_u f)^T(s) f(s) ds \right\| &\leq \left( \int_0^1 \frac{1}{12} \| (K_u f)(s) \|^2 ds \right)^{\frac{1}{2}} \left( \int_0^1 \frac{1}{12} \| f(s) \|^2 ds \right)^{\frac{1}{2}} \\ &\leq \frac{1}{12} \| \Gamma_u \| \int_0^1 \| f(s) \|^2 ds < \int_0^1 \| f(s) \|^2 ds \end{aligned}$$

and hence  $\| \frac{1}{12} K_u \| < 1$ . Thus

$$\det \left( I + \frac{1}{12} K_u \right) = \exp \left[ \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k} \text{tr} \left( \frac{1}{12} K_u \right)^k \right].$$

Therefore it suffices to show that for  $k \in \mathbb{N}$ ,

$$\text{tr} (K_u^k) = \text{tr} \int_{s_k=0}^1 \cdots \int_{s_1=0}^1 \left[ \prod_{l=1}^k \right] h(s_l, s_{l+1}) ds_1 \cdots ds_k = \gamma_k(u),$$



where by convention,  $s_{k+1} = s_1$ . Let

$$(K_u^k f)(s) = \int_0^1 p_k(s, s_1) f(s_1) ds_1,$$

where

$$p_1(s, s_1) = h(s, s_1)$$

and for  $k \geq 2$ ,

$$p_k(s, s_1) = \int_{s_k=0}^1 \cdots \int_{s_2=0}^1 h(s, s_2) \left[ \prod_{l=2}^k \right] h(s_l, s_{l+1}) ds_2 \cdots ds_k.$$

Let  $\Omega \subseteq L^2([0, 1])$  be an orthonormal basis and hence

$$\{\varpi e_i \mid \varpi \in \Omega, i = 1, 2, \dots, d\}$$

is an orthonormal basis for  $L^2([0, 1], \mathbb{R}^d)$ . Let  $(\cdot, \cdot)$  denote the inner product on  $L^2([0, 1], \mathbb{R}^d)$  and  $\langle \cdot, \cdot \rangle$  be the inner product on  $\mathbb{R}^d$ . Then

$$\begin{aligned} \text{tr}(K_u^k) &= \sum_{i, \varpi \in \Omega} (K_u^k \varpi e_i, \varpi e_i) = \sum_{i, \varpi \in \Omega} \int_0^1 \int_0^1 \langle p_k(s, t) \varpi(t) e_i, \varpi(s) e_i \rangle dt ds \\ &= \sum_{\varpi \in \Omega} \int_0^1 \int_0^1 \varpi(t) \varpi(s) \text{tr} p_k(s, t) dt ds \\ &= \int_0^1 \sum_{\varpi \in \Omega} \left( \int_0^1 \varpi(t) \text{tr} p_k(s, t) dt \right) \varpi(s) ds \\ &= \int_0^1 \text{tr} p_k(s, s) ds = \text{tr} \int_{s_k=0}^1 \cdots \int_{s_1=0}^1 \left[ \prod_{l=1}^k \right] h(s_l, s_{l+1}) ds_1 \cdots ds_k. \end{aligned}$$

Since  $K_u$  is trace class, all the interchanging between the integrals and the sums are valid. ■

### 5.3 Convergence of $\det(\widetilde{\mathcal{I}^n + \mathcal{X}^n}) \circ b_n$

**Lemma 5.27**

$$\det(\widetilde{\mathcal{I}^n + \mathcal{X}^n}) \circ b_n = 1_{\{n \geq n_0\}} \det(\mathcal{I}^n + \mathcal{X}^n) \circ b_n \longrightarrow 1$$

$\mu - a.s.$  as  $n \rightarrow \infty$ .

**Proof.** By Lemma 5.6, it suffices to consider  $W_\alpha(\mathbb{R}^d)$  with  $\frac{7}{15} < \alpha < \frac{1}{2}$ . Recall

$$\mathcal{X}^n = (\mathcal{I}^n + \mathcal{U}^n)^{-1} (\mathcal{S}^n)^T \mathcal{E}^n \mathcal{S}^n.$$

Now observe that  $\mathcal{I}^n + \mathcal{U}^n$  is a positive definite matrix with eigenvalues greater than or equal to 1. Thus since the norm of a symmetric matrix is equal to the maximum eigenvalue and the eigenvalues of  $(\mathcal{I}^n + \mathcal{U}^n)^{-1}$  is the reciprocal of the eigenvalues of  $\mathcal{I}^n + \mathcal{U}^n$ , we have

$$\|(\mathcal{I}^n + \mathcal{U}^n)^{-1}\| \leq 1.$$

Note that  $\|\mathcal{E}^n \circ b_n\| = O\left(\bigvee_{i=1,\dots,n} \|\Delta_i b\|^5\right)$  for  $n \geq n_0(\cdot)$ . Thus by Lemma 5.12, on  $W_\alpha(\mathbb{R}^d)$  and for  $n \geq n_0(\cdot)$ , we have

$$\|\mathcal{X}^n \circ b_n\| = O\left(n^2 \bigvee_{i=1,\dots,n} \|\Delta_i b\|^5\right).$$

Let  $\|\mathcal{X}^n \circ b_n\| \leq C[n^2 \bigvee_{i=1,\dots,n} \|\Delta_i b\|^5]$ ,  $C$  is a fixed constant independent of  $\omega$  and  $n$ . Therefore

$$\|\mathcal{X}^n \circ b_n\| \leq Cn^2 \bigvee_{i=1,\dots,n} \|\Delta_i b\|^5 < Cn^{2-5\alpha} < Cn^{-\frac{1}{3}}$$

and hence  $\|\mathcal{X}^n\| < 1$ . Thus using Equation (B.2), we have

$$\begin{aligned} \det[\mathcal{I}^n + \mathcal{X}^n] &= \exp\left[\sum_{k=1}^{\infty} \frac{1}{k} (-1)^{k+1} \text{Tr}[(\mathcal{X}^n)^k]\right] \\ &= \exp\left[\text{Tr} \mathcal{X}^n - \frac{1}{2} \text{Tr}[(\mathcal{X}^n)^2] + R_3(\mathcal{X}^n)\right], \end{aligned}$$

where  $R_3(\mathcal{X}^n) = \sum_{k=3}^{\infty} \frac{1}{k} (-1)^{k+1} \text{Tr}[(\mathcal{X}^n)^k]$ . Note that  $\mathcal{X}^n$  is a  $nd \times nd$  matrix and hence,

$$|R_3(\mathcal{X}^n \circ b_n)| \leq \frac{ndC^3 n^{3(2-5\alpha)}}{1 - Cn^{2-5\alpha}} = \frac{dC^3 n^{3(2-5\alpha)+1}}{1 - Cn^{2-5\alpha}}.$$

By the choice of  $\alpha$ ,  $\tau := 3(2 - 5\alpha) + 1 < 0$  and hence

$$|R_3(\mathcal{X}^n \circ b_n)| = O(n^\tau) \longrightarrow 0$$

as  $n \rightarrow \infty$ . Let  $P := (\mathcal{I}^n + \mathcal{U}^n \circ b_n)^{-1}$ ,  $S = \mathcal{S}^n$  and  $E = \mathcal{E}^n \circ b_n$  to simplify notation. Then, using Proposition C.1 twice,

$$\begin{aligned} |\text{Tr} \mathcal{X}^n \circ b_n| &= |\text{Tr}(PS^T ES)| = |\text{Tr}(SPS^T E)| \\ &\leq \|E\| \text{Tr}(SPS^T) = \|E\| \text{Tr}(S^T SP) \leq \|E\| \|P\| \text{Tr}(S^T S) \end{aligned}$$

where  $\text{Tr}(S^T S) = O(n^2)$ . This is because from Lemma 5.12, we have

$$\text{Tr}(S^T S) = \text{Tr} \mathcal{B}^n = \sum_{m=1}^n \text{mtr } I = \frac{dn(n+1)}{2}.$$

Thus

$$\begin{aligned}
|\text{Tr } \mathcal{X}^n \circ b_n| &\leq O(n^2) \|E\| = O(n^2(n^{-\alpha})^5) \\
&= O(n^2(n^{-7/15})^5) = O(n^{-1/3}) \\
&\longrightarrow 0
\end{aligned}$$

as  $n \rightarrow \infty$ . Similarly, using Lemma C.1 twice and the fact that  $P$  is positive definite so that  $SPS^T$  is positive definite, we have

$$\begin{aligned}
|\text{Tr } [(\mathcal{X}^n \circ b_n)^2]| &= |\text{Tr } (P^T E S P S^T E S)| = |\text{Tr } (S P S^T E S P S^T E)| \\
&\leq \text{Tr } (S P S^T) \|E S P S^T E\| = \text{Tr } (S^T S P) \|E S P S^T E\| \\
&\leq \text{Tr } (S^T S) \cdot \|P\| \|E S P S^T E\| \\
&\leq O(n^2) \|E S P S^T E\| \leq O(n^2) \|E\|^2 \|S\|^2 \\
&= O(n^4) \|E\|^2 = O(n^4(n^{-\alpha})^{10}) \\
&= O(n^4(n^{-7/15})^{10}) = O(n^4 n^{-14/3}) \\
&= O(n^{\frac{12-14}{3}}) = O(n^{-2/3}) \\
&\longrightarrow 0
\end{aligned}$$

as  $n \rightarrow \infty$ . Therefore,

$$\begin{aligned}
\det(\widetilde{\mathcal{I}^n + \mathcal{X}^n}) \circ b_n &= 1_{\{n \geq n_0\}} \det(\mathcal{I}^n + \mathcal{X}^n) \circ b_n \\
&= \exp \left[ \text{Tr } \mathcal{X}^n - \frac{1}{2} \text{Tr } [(\mathcal{X}^n)^2] + R_3(\mathcal{X}^n) \right] \circ b_n \\
&\longrightarrow 1
\end{aligned}$$

$\mu$ -a.s. as  $n \rightarrow \infty$ . ■

As a random variable on  $(W(\mathbb{R}^d), \mu)$ , the next theorem shows that  $\rho_n \circ \phi \circ b_n$  converges in  $\mu$ -measure.

**Theorem 5.28** *If  $\sup_{v \in O(M)} \|S_v\| < \frac{3}{17d}$ , then*

$$\rho_n \circ \phi \circ b_n \longrightarrow e^{-\frac{1}{6} \int_0^1 \text{Scal}(\tilde{\phi}(s)) \, ds} \sqrt{\det \left( I + \frac{1}{12} K_u \right)}.$$

in  $\mu$ -measure as  $n \rightarrow \infty$ .

**Proof.** It suffices to consider on  $W_\alpha(\mathbb{R}^d)$ , with  $\frac{7}{15} < \alpha < \frac{1}{2}$ . From Theorem 4.19,

$$\det[(\mathcal{F}^n)^T \mathcal{F}^n] = [\det(\mathcal{V}^n)]^2 \det(\mathcal{I}^n + \mathcal{U}^n) \det \left[ \mathcal{I}^n + \left( \mathcal{I}^n + \mathcal{U}^n \right)^{-1} (\mathcal{S}^n)^T \mathcal{E}^n \mathcal{S}^n \right].$$

Using Lemma 5.11, Theorem 5.23, Proposition 5.26 and Lemma 5.27, we have,

$$\begin{aligned} 1_{\{n \geq n_o\}} \det[\langle (\mathcal{F}^n)^T \mathcal{F}^n \rangle \circ b_n] &\longrightarrow e^{\frac{1}{3} \int_0^1 \text{Scal}(\tilde{\phi}(s)) \, ds + \gamma} \\ &= e^{\frac{1}{3} \int_0^1 \text{Scal}(\tilde{\phi}(s)) \, ds} \det \left( I + \frac{1}{12} K_u \right) \end{aligned}$$

in  $\mu$ -measure as  $n \rightarrow \infty$ . But by Theorem 4.7, for  $n \geq n_0(\cdot)$ ,

$$(\rho_n \circ \phi \circ b_n)^2 = \det[\langle (\mathcal{F}^n)^T \mathcal{F}^n \rangle] \circ b_n.$$

Thus taking the square root completes the proof. ■

## 6 $L^1$ Convergence of $\{\rho_n \circ \phi \circ b_n\}_{n=1}^\infty$

**Definition 6.1** Let  $\mu_{G_{\mathcal{P}}^1}$  be defined as in Theorem 1.4 on  $H_{\mathcal{P}}(\mathbb{R}^d)$  by the density

$$\frac{1}{Z_{\mathcal{P}}^1} e^{-\frac{1}{2} E_{\mathbb{R}^d}} \text{Vol}_{G_{\mathcal{P}}^1}$$

where  $Z_{\mathcal{P}}^1 := (2\pi)^{\frac{dn}{2}}$  and  $E_{\mathbb{R}^d}(\omega) := \int_0^1 \|\omega'(s)\|^2 \, ds$ .

The following 2 results can be found in [AD99] and hence we will omit the proofs.

**Theorem 6.2** Let  $\mu_{G_{\mathcal{P}}^1}$  be defined as in Definition 6.1. Write  $\phi_{\mathcal{P}} = \phi|_{H_{\mathcal{P}}(\mathbb{R}^d)}$ . Then  $\mu_{G_{\mathcal{P}}^1}$  is the pullback of  $\nu_{G_{\mathcal{P}}^1}$  by  $\phi_{\mathcal{P}}$ . i.e.

$$\mu_{G_{\mathcal{P}}^1} = (\phi_{\mathcal{P}})^* \nu_{G_{\mathcal{P}}^1}.$$

Let  $\pi_{\mathcal{P}} : W(\mathbb{R}^d) \rightarrow (\mathbb{R}^d)^n$  be given by

$$\pi_{\mathcal{P}}(\omega) := (\omega(s_1), \omega(s_2), \dots, \omega(s_n)).$$

Note that  $\pi_{\mathcal{P}} : H_{\mathcal{P}}(\mathbb{R}^d) \rightarrow (\mathbb{R}^d)^n$  is a linear isomorphism of finite dimensional vector spaces. Let  $i_{\mathcal{P}} : (\mathbb{R}^d)^n \rightarrow H_{\mathcal{P}}(\mathbb{R}^d)$  denote the inverse of  $\pi_{\mathcal{P}}|_{H_{\mathcal{P}}(\mathbb{R}^d)}$ .

**Lemma 6.3** Let  $(y_1, \dots, y_n)$  denote the standard cartesian coordinates on  $(\mathbb{R}^d)^n$  and  $y_0 := 0$ . Then

$$i_{\mathcal{P}}^* \mu_{G_{\mathcal{P}}^1} = \frac{1}{Z_{\mathcal{P}}^1} \left( \prod_{i=1}^n (\Delta_i s)^{-\frac{d}{2}} \exp \left( -\frac{1}{2\Delta_i s} \|y_i - y_{i-1}\|^2 \right) \right) dy_1 dy_2 \dots dy_n. \quad (6.1)$$

This equation can also be written as

$$i_{\mathcal{P}}^* \mu_{G_{\mathcal{P}}^1} = \left( \prod_{i=1}^n p_{\Delta_i s}(y_{i-1}, y_i) \right) dy_1 dy_2 \dots dy_n$$

where  $p_s(x, y) := (\frac{1}{2\pi})^{-\frac{d}{2}} \exp(-\frac{\|x-y\|^2}{2s})$  is the heat kernel on  $\mathbb{R}^d$ .

We are now ready to prove the main result.

**Theorem 6.4** *Let  $M$  be a compact Riemannian manifold,  $f : W(M) \rightarrow \mathbb{R}$  be a bounded continuous function and  $\mathcal{P} = \{0 < \frac{1}{n} < \dots < \frac{n}{n} = 1\}$  is an equally spaced partition. Let  $b$  be brownian motion,  $u$  solves Equation (5.1),  $\tilde{\phi} = \pi \circ u$  and  $\widetilde{\llcorner}$  is stochastic parallel translation defined in Definition 1.12. Suppose  $\sup_{v \in O(M)} \|S_v\| < \frac{3}{17d}$ , where  $S_v$  is sectional curvature and  $\rho_{\mathcal{P}} \circ \phi \circ b_{\mathcal{P}}$  is uniformly integrable. Then*

$$\begin{aligned} \int_{H_{\mathcal{P}}(M)} f(\sigma) d\nu_{\mathcal{P}}(\sigma) &= \int_{H_{\mathcal{P}}(M)} f(\sigma) \rho_{\mathcal{P}}(\sigma) d\nu_{G_{\mathcal{P}}^1}(\sigma) \\ &\xrightarrow{|\mathcal{P}| \rightarrow 0} \int_{W(M)} f(\sigma) e^{-\frac{1}{6} \int_0^1 \text{Scal}(\sigma(s)) ds} \sqrt{\det \left( I + \frac{1}{12} K_{\widetilde{\llcorner}(\sigma)} \right)} d\nu(\sigma) \end{aligned}$$

where  $\nu$  is Wiener measure on  $M$ .  $\text{Scal}$  is the scalar curvature of the manifold and  $K_{\widetilde{\llcorner}(\sigma)} : L^2([0, 1] \rightarrow \mathbb{R}^d) \longrightarrow L^2([0, 1] \rightarrow \mathbb{R}^d)$  is given by

$$\left( K_{\widetilde{\llcorner}(\sigma)} v \right) (s) = \int_0^1 (s \wedge t) \Gamma(\widetilde{\llcorner}_t(\sigma)) v(t) dt$$

where

$$\Gamma(\widetilde{\llcorner}_t(\sigma)) = \sum_{i,j=1}^d \left\{ \begin{aligned} &\Omega_{\widetilde{\llcorner}_t(\sigma)}(e_i, \Omega_{\widetilde{\llcorner}_t(\sigma)}(e_i, \cdot) e_j) e_j + \Omega_{\widetilde{\llcorner}_t(\sigma)}(e_i, \Omega_{\widetilde{\llcorner}_t(\sigma)}(e_j, \cdot) e_i) e_j \\ &+ \Omega_{\widetilde{\llcorner}_t(\sigma)}(e_i, \Omega_{\widetilde{\llcorner}_t(\sigma)}(e_j, \cdot) e_i) e_i \end{aligned} \right\}$$

for any orthonormal basis  $\{e_i\}_{i=1}^d \subseteq T_o M$ , and

$$\Omega_{\widetilde{\llcorner}_t(\sigma)}(a, c) := \widetilde{\llcorner}_t^{-1}(\sigma) R \left( \widetilde{\llcorner}_t(\sigma) a, \widetilde{\llcorner}_t(\sigma) c \right) \widetilde{\llcorner}_t(\sigma)$$

for any vectors  $a, c \in T_o M$ ,  $R$  being the curvature tensor.

**Proof.** By Theorem 5.2,  $f \circ \phi \circ b_{\mathcal{P}} := f_{\mathcal{P}}$  converges to  $f \circ \tilde{\phi} \mu - a.s.$  as  $|\mathcal{P}| \rightarrow 0$ . Thus we have

$$\begin{aligned} \int_{H_{\mathcal{P}}(M)} f d\nu_{\mathcal{P}} &= \int_{H_{\mathcal{P}}(M)} f \rho_{\mathcal{P}} d\nu_{G_{\mathcal{P}}^1} \\ &= \int_{H_{\mathcal{P}}(\mathbb{R}^d)} (f \rho_{\mathcal{P}}) \circ \phi d\mu_{G_{\mathcal{P}}^1} \text{ (By Theorem 6.2)} \\ &= \int_{W(\mathbb{R}^d)} (f \rho_{\mathcal{P}}) \circ \phi \circ b_{\mathcal{P}} d\mu \text{ (By Lemma 6.3)} \\ &= \int_{W(\mathbb{R}^d)} f_{\mathcal{P}} \cdot \rho_{\mathcal{P}} \circ \phi \circ b_{\mathcal{P}} d\mu. \end{aligned}$$

By the assumption on  $\rho_{\mathcal{P}} \circ \phi \circ b_{\mathcal{P}}$ , since  $f$  is bounded and continuous, we have  $f_{\mathcal{P}} \cdot \rho_{\mathcal{P}} \circ \phi \circ b_{\mathcal{P}}$  is uniformly integrable. Therefore by Theorem 5.28,

$$\begin{aligned} \int_{H_{\mathcal{P}}(M)} f(\sigma) d\nu_{\mathcal{P}}(\sigma) &= \int_{W(\mathbb{R}^d)} (f_{\mathcal{P}} \cdot \rho_{\mathcal{P}} \circ \phi \circ b_{\mathcal{P}})(\omega) d\mu(\omega) \\ &\longrightarrow \int_{W(\mathbb{R}^d)} \left( f \circ \tilde{\phi} \cdot e^{-\frac{1}{6} \int_0^1 \text{Scal}(\tilde{\phi}(s)) ds} \sqrt{\det \left( I + \frac{1}{12} K_u \right)} \right)(\omega) d\mu(\omega) \\ &= \int_{W(M)} f(\sigma) e^{-\frac{1}{6} \int_0^1 \text{Scal}(\sigma(s)) ds} \sqrt{\det \left( I + \frac{1}{12} K_{\tilde{\gamma}(\sigma)} \right)} d\nu(\sigma) \end{aligned}$$

as  $|\mathcal{P}| \rightarrow 0$ , where  $K_u$  was defined in Definition 5.24. Note that  $\nu = \mu \tilde{\phi}^{-1}$  and  $u = \tilde{\gamma}(\tilde{\phi})$  from (1) and (2) of Remark 5.4 respectively.  $\blacksquare$

**Corollary 6.5** *Let  $M$  be a compact Riemannian manifold,  $f : W(M) \rightarrow \mathbb{R}$  be a bounded continuous function and  $\mathcal{P}$  is an equally spaced partition. Suppose that  $\forall v \in O(M)$ ,  $0 \leq S_v < \frac{3}{17d}$ . Then*

$$\begin{aligned} \int_{H_{\mathcal{P}}(M)} f(\sigma) d\nu_{\mathcal{P}}(\sigma) \\ \xrightarrow{|\mathcal{P}| \rightarrow 0} \int_{W(M)} f(\sigma) e^{-\frac{1}{6} \int_0^1 \text{Scal}(\sigma(s)) ds} \sqrt{\det \left( I + \frac{1}{12} K_{\tilde{\gamma}(\sigma)} \right)} d\nu(\sigma). \end{aligned}$$

**Proof.** By Theorem 3.15, we know that  $\rho_{\mathcal{P}} \circ \phi \circ b_{\mathcal{P}}$  is uniformly integrable under the assumptions on the sectional curvature. Hence the corollary now follows from Theorem 6.4.  $\blacksquare$

## 7 Further Questions

Two questions immediately arise from Corollary 6.5.

1. Can we remove the upper bound restriction on the sectional curvature? In other words, does the result hold true for any compact manifold with non-negative sectional curvature?
2. Can the result be extended to an arbitrary compact manifold?

The restriction of the result on a non-negative manifold arises when we are trying to prove the uniform integrability of  $\rho_{\mathcal{P}}$ . Thus one way to improve the result is to obtain a better upper estimate for  $\rho_{\mathcal{P}}$ , and show that without any restrictions on the sectional curvature of the manifold,  $\rho_{\mathcal{P}}$  is still uniformly integrable.

One possible research problem will be to consider a different metric on  $H(M)$ . For example, by considering a  $L^2$ -metric  $G^0$  on  $TH(M)$  where

$$G^0(X, X) := \int_0^1 g(X(s), X(s)) \, ds,$$

for  $X \in T_\sigma H(M)$ . Instead of considering Wiener space  $W(M)$ , one can consider the space of pinned paths, equipped with pinned Wiener measure and carry out a similar analysis.

## 8 Acknowledgements

The author would like to thank Professor Bruce Driver for all his help and input into this paper.

## A Trace Class Operators

A background knowledge on trace class operators can be found in [DS88b] and [Sim79].

In general, it is difficult to determine if an integral operator is trace class. However, the following theorem, taken from Theorem 2.12 of [Sim79], gives a condition for which an integral operator is trace class. See also Section XI.4 of [RS79].

**Theorem A.1** *Let  $\mu$  be a Baire measure on a locally compact space  $X$ . Let  $K$  be a function on  $X \times X$  which is continuous and Hermitian positive, that is*

$$\sum_{i,j=1}^N \overline{z_i} z_j K(x_i, x_j) \geq 0$$

*for any  $x_1, \dots, x_N \in X$ ,  $z_1, \dots, z_N \in \mathbb{C}^N$  and for any  $N$ . Then  $K(x, x) \geq 0$  for all  $x$ . Suppose that in addition,*

$$\int K(x, x) \, d\mu(x) < \infty.$$

*Then there exists a unique trace class integral operator  $A$  such that*

$$(Af)(x) = \int K(x, y) f(y) \, d\mu(y)$$

*and*

$$\|A\|_1 = \int K(x, x) \, d\mu.$$

**Proposition A.2** *Let  $(Af)(s) = \int_0^1 (s \wedge t) f(t) \, dt$ . Then  $A$  is a trace class operator.*

**Proof.** Let  $X = [0, 1]$  and  $\mu$  be Lebesgue measure. Using Theorem A.1, it suffices to show that  $s \wedge t$  is Hermitian positive. Let  $z_1, z_2, \dots, z_N$  be any complex numbers and let  $x_1, \dots, x_N \in [0, 1]$ . The proof is by induction. Clearly when  $N = 1$ , it is trivial. Suppose it is true for all values from  $k = 1, 2, \dots, N - 1$ . By relabelling, we can assume that  $x_1 \leq x_k$ ,  $k = 2, \dots, N$ . Hence  $x_1 \wedge x_k = x_1$  for any  $k$ . Thus

$$\begin{aligned} \sum_{j=1}^N \overline{z_1} z_j (x_1 \wedge x_j) + \sum_{j=1}^N \overline{z_j} z_1 (x_j \wedge x_1) &= x_1 \sum_{j=1}^N \overline{z_1} z_j + x_1 \sum_{j=1}^N \overline{z_j} z_1 \\ &= x_1 \left( \sum_{j=1}^N \overline{z_j} \right) \left( \sum_{j=1}^N z_j \right) - x_1 \sum_{i,j=2}^N \overline{z_i} z_j. \end{aligned}$$

Therefore,

$$\begin{aligned} \sum_{i,j=1}^N \overline{z_i} z_j (x_i \wedge x_j) &= x_1 \left( \sum_{j=1}^N \overline{z_j} \right) \left( \sum_{j=1}^N z_j \right) - x_1 \sum_{i,j=2}^N \overline{z_i} z_j + \sum_{i,j=2}^N \overline{z_i} z_j (x_i \wedge x_j) \\ &= x_1 \left( \sum_{j=1}^N \overline{z_j} \right) \left( \sum_{j=1}^N z_j \right) + \sum_{i,j=2}^N \overline{z_i} z_j (c_i \wedge c_j), \end{aligned}$$

where  $c_i = x_i - x_1 \geq 0$ ,  $i = 2, \dots, N$ . Thus by induction hypothesis,

$$\sum_{i,j=2}^N \overline{z_i} z_j (c_i \wedge c_j) \geq 0$$

and hence

$$\sum_{i,j=1}^N \overline{z_i} z_j (x_i \wedge x_j) \geq 0.$$

■

**Proposition A.3** *Let  $\mathcal{H}$  be a separable Hilbert space. Suppose  $A$  is trace class and  $B$  is a bounded operator. Then*

$$\operatorname{tr} |AB| \leq \|B\| \operatorname{tr} |A|$$

where  $\|\cdot\|$  is the operator norm.

For a proof, see Theorem VI.25 in [RS80].

## B Perturbation Formulas

**Lemma B.1** *Let  $U$  be a  $d \times d$  matrix and for  $r \in \mathbb{N}$ , let*

$$\Psi_r(U) := \sum_{k=1}^r (-1)^{k+1} \frac{1}{k} \operatorname{tr} U^k.$$



If  $U$  is positive definite, then there exists  $R_{r+1}(U)$  such that

$$\det(I + U) = \exp(\Psi_r(U) + R_{r+1}(U)), \quad (\text{B.1})$$

where

$$|R_{r+1}(U)| \leq \frac{1}{r+1} \text{tr } U^{r+1}.$$

If  $U$  is any  $d \times d$  matrix (not necessarily positive) such that  $\|U\| < 1$ , then

$$\det(I + U) = \exp(\Psi_r(U) + R_{r+1}(U)), \quad (\text{B.2})$$

with

$$R_{r+1}(U) = \sum_{k=r+1}^{\infty} (-1)^{k+1} \frac{1}{k} \text{tr } U^k$$

such that

$$|R_{r+1}(U)| \leq \frac{d \|U\|^{r+1}}{1 - \|U\|}. \quad (\text{B.3})$$

**Proof.** Let  $\{\lambda_1, \lambda_2, \dots, \lambda_d\}$  be the set of eigenvalues of  $U$ . Then

$$\begin{aligned} \det(I + U) &= \prod_{i=1}^d (1 + \lambda_i) \\ &= \exp \left( \sum_{i=1}^d \ln(1 + \lambda_i) \right), \end{aligned} \quad (\text{B.4})$$

since by assumption,  $\lambda_i \geq 0$  for all  $i$ . But by applying Taylor's Theorem to  $\ln(1 + x)$ , we have

$$\ln(1 + x) = \sum_{k=1}^r (-1)^{k+1} \frac{1}{k} x^k + R_{r+1}(x),$$

where

$$R_{r+1}(x) = x^{r+1} \frac{(-1)^r}{r+1} \frac{1}{(1+c)^{r+1}}$$

for some  $c \in (0, x)$ . Therefore,

$$|R_{r+1}(x)| \leq \frac{1}{r+1} x^{r+1}.$$

Since

$$\text{tr } U^k = \sum_{i=1}^d \lambda_i^k,$$

we have

$$\begin{aligned} \sum_{i=1}^d \ln(1 + \lambda_i) &= \sum_{i=1}^d \left( \sum_{k=1}^r (-1)^{k+1} \frac{1}{k} \lambda_i^k + R_{r+1}(\lambda_i) \right) \\ &= \sum_{k=1}^r (-1)^{k+1} \frac{1}{k} \text{tr } U^k + \sum_{i=1}^d R_{r+1}(\lambda_i) \\ &= \Psi_r(U) + R_{r+1}(U), \end{aligned}$$

where  $R_{r+1}(U) := \sum_{i=1}^d R_{r+1}(\lambda_i)$  and  $|R_{r+1}(U)| \leq \sum_{i=1}^d \lambda_i^{r+1} = \frac{1}{r+1} \text{tr } U^{r+1}$ . To prove Equation (B.2), we assume that  $\|U\| < 1$ . Thus the eigenvalues of  $U$ ,  $\lambda_i < 1$  for all  $i$ . Therefore Equation (B.4) holds. But  $\ln(1 + \lambda_i) = \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k} \lambda_i^k$  and this sum converges absolutely. Thus

$$\begin{aligned} \sum_{i=1}^d \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k} \lambda_i^k &= \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k} \sum_{i=1}^d \lambda_i^k \\ &= \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k} \text{tr } U^k. \end{aligned}$$

Hence this proves Equation (B.2). In this case,

$$\begin{aligned} |R_{r+1}(U)| &= \left| \sum_{k=r+1}^{\infty} (-1)^{k+1} \frac{1}{k} \text{tr } U^k \right| \leq \sum_{k=r+1}^{\infty} \frac{1}{k} |\text{tr } U^k| \\ &\leq \sum_{k=r+1}^{\infty} d \|U\|^k = \frac{d \|U\|^{r+1}}{1 - \|U\|}. \end{aligned} \tag{B.5}$$

Alternatively, Equation (B.2) can be proved by a rewriting of the standard formula,

$$\begin{aligned} \frac{d}{ds} \log(\det(I - sU)) &= -\text{tr}((I - sU)^{-1}U) \\ &= -\text{tr} \left( \sum_{k=0}^{\infty} s^k U^k U \right) \\ &= -\sum_{k=0}^{\infty} s^k \text{tr } (U^{k+1}). \end{aligned}$$

■

Let  $\mathcal{J}_1$  be the space of trace class operators, equipped with the norm  $\|\cdot\|_1$ , define by

$$\|A\|_1 = \text{tr } |A|.$$

Let  $A \in \mathcal{J}_1$ . Suppose  $-z^{-1}$  lies in the resolvent set of  $A$ , i.e.  $-z^{-1} \notin \sigma(A)$ . Then the mapping  $A \rightarrow \ln(I + zA)$  is defined and is continuous on  $A$ . (See Lemma 15 in Section XI.9.22 of [DS88b].) Thus, one can define a function  $\det(I + \cdot)$  on  $\mathcal{J}_1$  in the following way.

**Definition B.2** Define for  $-z^{-1} \notin \sigma(A)$ ,

$$\det(I + zA) = \exp(\text{tr } \ln(I + zA)).$$

This function  $\det(I + zA)$  is analytic in  $z$  and has only removable singularities at the points  $z$  such that  $-z^{-1} \in \sigma(T)$ . We can now show that Equation (B.2) holds even for trace class operator.

**Lemma B.3** Let  $A$  be a trace class operator on a Hilbert space  $\mathcal{H}$ , with  $\|A\| < 1$  where  $\|\cdot\|$  is the operator norm. Then Equation (B.2) holds with  $U$  replaced by  $A$ .

**Proof.** Since  $\text{tr}$  is a continuous function on  $\mathcal{J}_1$  (See Theorem 19 in Section XI.9.20 of [DS88b].), we have

$$\begin{aligned} \text{tr} \ln(I + zA) &= \text{tr} \lim_{N \rightarrow \infty} \sum_{k=1}^N \frac{(-1)^{k+1}}{k} (zA)^k = \lim_{N \rightarrow \infty} \text{tr} \sum_{k=1}^N \frac{(-1)^{k+1}}{k} (zA)^k \\ &= \lim_{N \rightarrow \infty} \sum_{k=1}^N \frac{(-1)^{k+1}}{k} \text{tr} (zA)^k = \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} \text{tr} (zA)^k. \end{aligned}$$

Because  $\|A\| < 1$ , setting  $z = 1$  gives us our desired result. ■

## C Matrix Inequalities

**Proposition C.1** If  $A, B$  are two  $N \times N$  matrices with  $B$  being positive semi definite, then

$$|\text{tr}(AB)| \leq \|A\| \text{tr} B \quad (\text{C.1})$$

and in particular by taking  $B = I$ ,

$$|\text{tr} A| \leq N \|A\|. \quad (\text{C.2})$$

**Proof.** Let  $\{e_i\}_{i=1,2,\dots,N}$  be an orthonormal basis of eigenvectors of  $B$  with corresponding eigenvalues,  $\{\lambda_i \geq 0\}_{i=1,2,\dots,N}$ . Then

$$\begin{aligned} |\text{tr}(AB)| &= \left| \sum_{i=1}^N \langle AB e_i, e_i \rangle \right| = \left| \sum_{i=1}^N \lambda_i \langle A e_i, e_i \rangle \right| \\ &\leq \sum_{i=1}^N \lambda_i |\langle A e_i, e_i \rangle| \leq \sum_{i=1}^N \lambda_i \|A\| = \|A\| \text{tr} B. \end{aligned}$$

■

**Proposition C.2** Suppose that  $M$  is a positive definite  $N \times N$  matrix and  $\alpha > 0$ . Then

$$\det(M) \leq \left( \frac{\text{tr}(M)}{N} \right)^N \leq \alpha^N e^{\text{tr}(\alpha^{-1}M - I)}. \quad (\text{C.3})$$

Moreover if  $\alpha \geq 1$ , then

$$\det(M) \leq \alpha^N e^{\alpha^{-1} \text{tr}(M - I)}. \quad (\text{C.4})$$

**Proof.** Let  $\{\lambda_i\}_{i=1}^N$  be the eigenvalues of  $M$ , then

$$\det M = \lambda_1 \cdots \lambda_N = \left( \lambda_1^{1/N} \cdots \lambda_N^{1/N} \right)^N \leq \left( \sum_{i=1}^N \frac{1}{N} \lambda_i \right)^N = \left( \frac{\text{tr}(M)}{N} \right)^N.$$

Now suppose that  $\alpha > 0$ . Then

$$\begin{aligned} \det M &= \alpha^N \det(\alpha^{-1}M) \leq \alpha^N \left( \frac{\text{tr}(\alpha^{-1}M)}{N} \right)^N \\ &\leq \alpha^N \left( 1 + \frac{\text{tr}(\alpha^{-1}M - I)}{N} \right)^N \leq \alpha^N e^{\text{tr}(\alpha^{-1}M - I)}, \end{aligned}$$

where the last inequality follows from the inequality

$$\left( 1 + \frac{x}{N} \right)^N \leq e^x \text{ for all } N \in \mathbb{N} \text{ and } x \geq -N.$$

Indeed, elementary calculus shows that  $f(x) := e^{-x} \left( 1 + \frac{x}{N} \right)^N$  for  $x \geq -N$  has a global maximum of one at  $x = 0$ .

Alternatively,

$$\begin{aligned} \det(M) &= \alpha^N \det(\alpha^{-1}M) = \alpha^N \det(\alpha^{-1}M - I + I) \\ &= \alpha^N \prod_{i=1}^N (1 + (\alpha^{-1}\lambda_i - 1)) \leq \alpha^N \prod_{i=1}^N e^{(\alpha^{-1}\lambda_i - 1)} = \alpha^N e^{\text{tr}(\alpha^{-1}M - I)} \end{aligned}$$

wherein we have used the inequality,  $1 + x \leq e^x$ , which results from the convexity of the exponential function. Finally, by optimizing this inequality over  $\alpha > 0$ , (take  $\alpha = N^{-1} \text{tr} M$ ), the previous inequality implies  $\det M \leq (N^{-1} \text{tr} M)^N$ . ■

## D Bounds on Curvature

**Definition D.1** Let  $M$  be a manifold,  $u \in O(M)$  and  $\Omega_u$  as defined in Definition 2.1. Define for any vectors  $x, y, w, z \in T_o M$ ,

$$R_u(x, y, w, z) = \langle \Omega_u(x, y)w, z \rangle,$$

and for any linearly independent vectors  $x, y$ ,

$$S_u(x, y) = \frac{R_u(x, y, x, y)}{\|x\|^2 \|y\|^2 - \langle x, y \rangle^2}.$$

The curvature tensor,  $R_u$  satisfies the following symmetries,

$$R_u(x, y, w, z) = -R_u(y, x, w, z), \quad (\text{D.1})$$

$$R_u(x, y, w, z) = -R_u(x, y, z, w), \quad (\text{D.2})$$

$$R_u(x, y, w, z) = R_u(w, z, x, y), \quad (\text{D.3})$$

$$R_u(x, y, w, z) + R_u(w, x, y, z) + R_u(y, w, x, z) = 0. \quad (\text{D.4})$$

The last equality is the Bianchi Identity.

$S_u$  is called the sectional curvature and if we define  $S_u(V) := S_u(x, y)$  where  $V$  is the plane spanned by any linearly independent vectors  $x, y$ , then it can be shown that  $S_u(V)$  is independent of the choice of  $x$  and  $y$ . We will write

$$\| S_u \| = \sup\{|S_u(V)| \mid V \text{ is a plane in } T_0M\},$$

and

$$K = \sup_{u \in O(M)} \| S_u \|.$$

Since we are considering a compact manifold  $M$ ,  $K$  is finite. Immediate from this definition, we have

$$|R_u(x, y, x, y)| \leq \| S_u \| \| x \|^2 \| y \|^2 \quad (\text{D.5})$$

for any  $x, y \in T_oM$ .

For any vector  $v$ , Equation (D.3) shows that  $\Omega_u(v, \cdot)v$  is a symmetric matrix. It follows that

$$\| \Omega_u(v, \cdot)v \| = \sup_{\|w\|=1} \| \Omega_u(v, w)v \| = \sup_{\|w\|=1} |\langle \Omega_u(v, w)v, w \rangle| \leq \| S_u \| \| v \|^2.$$

The last inequality follows from Equation (D.5). As a consequence, if we take the supremum over  $O(M)$ , then

$$\sup_{u \in O(M)} \| \Omega_u(v, \cdot)v \| \leq K \| v \|^2.$$

**Proposition D.2** *If  $\sup_{u \in O(M)} \| S_u \| < \frac{3}{17d}$ , then for any orthonormal frame  $\{e_i\}_{i=1}^d \subseteq T_oM$ ,*

$$\sup_{u \in O(M)} \| \Omega_u(e_i, \cdot)e_j \| < \frac{2}{d},$$

where  $d$  is the dimension of the manifold.

**Proof.** Let  $x, y, w, z \in T_o M$ . Then using Equations (D.1), (D.2) and (D.3),

$$\begin{aligned}
A &:= R_u(x+w, y+z, x+w, y+z) - R_u(x+w, y, x+w, y) - R_u(x+w, z, x+w, z) \\
&\quad - R_u(x, y+z, x, y+z) - R_u(w, y+z, w, y+z) + R_u(w, y, w, y) + R_u(x, z, x, z) \\
&= R_u(x+w, y, x+w, z) + R_u(x+w, z, x+w, y) - R_u(x, y, x, z) - R_u(x, z, x, y) - R_u(x, y, x, y) \\
&\quad - R_u(w, y, w, z) - R_u(w, z, w, y) - R_u(w, z, w, z) \\
&= R_u(x, y, w, z) + R_u(w, y, x, z) + R_u(x, z, w, y) + R_u(w, z, x, y) - R_u(x, y, x, y) - R_u(w, z, w, z) \\
&= 2R_u(x, y, w, z) + 2R_u(w, y, x, z) - R_u(x, y, x, y) - R_u(w, z, w, z).
\end{aligned}$$

Similarly,

$$\begin{aligned}
B &:= R_u(x+z, y+w, x+z, y+w) - R_u(x+z, y, x+z, y) - R_u(x+z, w, x+z, w) \\
&\quad - R_u(x, y+w, x, y+w) - R_u(z, y+w, z, y+w) + R_u(x, w, x, w) + R_u(z, y, z, y) \\
&= R_u(x+z, y, x+z, w) + R_u(x+z, w, x+z, y) - R_u(x, y, x, w) - R_u(x, w, x, y) - R_u(x, y, x, y) \\
&\quad - R_u(z, y, z, w) - R_u(z, w, z, y) - R_u(z, w, z, w) \\
&= R_u(x, y, z, w) + R_u(z, y, x, w) + R_u(x, w, z, y) + R_u(z, w, x, y) - R_u(x, y, x, y) - R_u(z, w, z, w) \\
&= 2R_u(x, y, z, w) + 2R_u(z, y, x, w) - R_u(x, y, x, y) - R_u(w, z, w, z).
\end{aligned}$$

Thus

$$\begin{aligned}
A - B &= 2R_u(x, y, w, z) + 2R_u(w, y, x, z) - 2R_u(x, y, z, w) - 2R_u(z, y, x, w) \\
&= 2R_u(x, y, w, z) + 2R_u(x, y, w, z) + 2R_u(w, y, x, z) + 2R_u(y, z, x, w) \\
&= 4R_u(x, y, w, z) + 2R_u(w, y, x, z) + 2R_u(x, w, y, z) \\
&= 4R_u(x, y, w, z) - 2R_u(y, x, w, z) \\
&= 6R_u(x, y, w, z).
\end{aligned}$$

The second last equality follows from the Bianchi Identity. Hence for any unit vectors  $x, y, w, z$ ,

$$\begin{aligned}
|A| &\leq \|S_u\| [\|x+w\|^2 \|y+z\|^2 + 2\|x+w\|^2 + 2\|y+z\|^2 + 2] \\
&\leq \|S_u\| [4^2 + 2 \cdot 8 + 2] = 34 \|S_u\|.
\end{aligned}$$

Similarly,  $|B| \leq 34 \|S_u\|$ . Therefore, for any unit vectors  $x, y, w, z$ ,

$$|R_u(x, y, w, z)| = \frac{1}{6} |A - B| \leq \frac{34}{3} \|S_u\|.$$

and hence

$$\sup_{\|x\|=\|y\|=\|w\|=\|z\|=1} |R_u(x, z, w, z)| \leq \frac{34}{3} \|S_u\|.$$

Now for any  $u \in O(M)$  and orthonormal frame  $\{e_i\}_{i=1}^d$ , let  $\Omega_{ij}(y) := \Omega_u(e_i, y)e_j$ . Then

$$\begin{aligned} \sup_{\|y\|=1} \|\Omega_{ij}(y)\| &= \sup_{\|y\|=1} \left\langle \Omega_{ij}(y), \frac{\Omega_{ij}(y)}{\|\Omega_{ij}(y)\|} \right\rangle = \sup_{\|y\|=1} \left| R_u \left( e_i, y, e_j, \frac{\Omega_{ij}(y)}{\|\Omega_{ij}(y)\|} \right) \right| \\ &\leq \sup_{\|x\|=\|y\|=\|w\|=\|z\|=1} |R_u(x, y, w, z)| \leq \frac{34}{3} \|S_u\|. \end{aligned}$$

So, for any  $u \in O(M)$  and orthonormal basis  $\{e_i\}_{i=1}^d \subseteq T_o M$ ,  $\|\Omega_{ij}\| \leq \frac{34}{3} \|S_u\|$ . If we choose  $\sup_{u \in O(M)} \|S_u\| < 3/(17d)$ , then

$$\sup_{u \in O(M)} \|\Omega_u(e_i, \cdot)e_j\| < \frac{34}{3} \cdot \frac{3}{17d} = \frac{2}{d}.$$

■

## References

- [AD99] Lars Andersson and Bruce K. Driver, *Finite-dimensional approximations to Wiener measure and path integral formulas on manifolds*, J. Funct. Anal. **165** (1999), no. 2, 430–498. MR MR1698956 (2000j:58059)
- [Bis84] Jean-Michel Bismut, *Large deviations and the Malliavin calculus*, Progress in Mathematics, vol. 45, Birkhäuser Boston Inc., Boston, MA, 1984. MR MR755001 (86f:58150)
- [Che72] K.S. Cheng, *Quantization of a general dynamical system by feynman's path integration formulation*, Journal of Mathematical Physics **13** (1972), no. 11, 1723–1726.
- [CM96] Ana-Bela Cruzeiro and Paul Malliavin, *Renormalized differential geometry on path space: structural equation, curvature*, J. Funct. Anal. **139** (1996), no. 1, 119–181. MR MR1399688 (97h:58175)
- [CMHM78] Alexandre J. Chorin, Marjorie F. McCracken, Thomas J. R. Hughes, and Jerrold E. Marsden, *Product formulas and numerical algorithms*, Comm. Pure Appl. Math. **31** (1978), no. 2, 205–256. MR MR0488713 (58 #8230)
- [Dar84] R. W. R. Darling, *On the convergence of Gangolli processes to Brownian motion on a manifold*, Stochastics **12** (1984), no. 3-4, 277–301. MR MR749378 (86i:58139)
- [dC92] Manfredo Perdigão do Carmo, *Riemannian geometry*, Mathematics: Theory & Applications, Birkhäuser Boston Inc., Boston, MA, 1992, Translated from the second Portuguese edition by Francis Flaherty. MR MR1138207 (92i:53001)

- [DeW72] Cécile Morette DeWitt, *Feynman's path integral*, Comm. Math. Phys. **28** (1972), 47–67, Definition without limiting procedure. MR MR0309456 (46 #8565)
- [DeW92] Bryce DeWitt, *Supermanifolds*, second ed., Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 1992. MR MR1172996 (93j:58008)
- [DH98] B. K. Driver and Y. Hu, *Wong zakai type approximation theorems*, <http://math.ucsd.edu/~driver/>, 1998.
- [Dir33] P. A. M. Dirac, Physikalische Zeitschrift der Sowjetunion **3** (1933), 64.
- [DM75] Cecile DeWitt-Morette, *Path integrals in Riemannian manifolds*, International Symposium on Mathematical Problems in Theoretical Physics (Kyoto Univ., Kyoto, 1975), Springer, Berlin, 1975, pp. 535–542. Lecture Notes in Phys., 39. MR MR0674340 (58 #32618)
- [DME81] Cécile DeWitt-Morette and K. D. Elworthy (eds.), *New stochastic methods in physics*, North-Holland Publishing Co., Amsterdam, 1981, Phys. Rep. **77** (1981), no. 3. MR MR639022 (83b:81009)
- [DMENS80] C. DeWitt-Morette, K. D. Elworthy, B. L. Nelson, and G. S. Sammelman, *A stochastic scheme for constructing solutions of the Schrödinger equations*, Ann. Inst. H. Poincaré Sect. A (N.S.) **32** (1980), no. 4, 327–341. MR MR594632 (82m:81028)
- [DMMN79] Cécile DeWitt-Morette, Amar Maheshwari, and Bruce Nelson, *Path integration in nonrelativistic quantum mechanics*, Phys. Rep. **50** (1979), no. 5, 255–372. MR MR525283 (80k:81039)
- [Dri89a] Bruce K. Driver, *Classifications of bundle connection pairs by parallel translation and lassos*, J. Funct. Anal. **83** (1989), no. 1, 185–231. MR MR993446 (90b:58033)
- [Dri89b] ———, *YM<sub>2</sub>: continuum expectations, lattice convergence, and lassos*, Comm. Math. Phys. **123** (1989), no. 4, 575–616. MR MR1006295 (91c:81049)
- [Dri92] ———, *A Cameron-Martin type quasi-invariance theorem for Brownian motion on a compact Riemannian manifold*, J. Funct. Anal. **110** (1992), no. 2, 272–376. MR MR1194990 (94a:58214)
- [DS88a] Nelson Dunford and Jacob T. Schwartz, *Linear operators. Part I*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1988, General theory, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1958 original, A Wiley-Interscience Publication. MR MR1009162 (90g:47001a)



- [DS88b] ———, *Linear operators. Part II*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1988, Spectral theory. Selfadjoint operators in Hilbert space, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1963 original, A Wiley-Interscience Publication. MR MR1009163 (90g:47001b)
- [EE71] J. Eells and K. D. Elworthy, *Wiener integration on certain manifolds*, Problems in non-linear analysis (C.I.M.E., IV Ciclo, Varenna, 1970), Edizioni Cremonese, Rome, 1971, pp. 67–94. MR MR0346835 (49 #11557)
- [Eel58] James Eells, Jr., *On the geometry of function spaces*, Symposium internacional de topología algebraica International symposium on algebraic topology, Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958, pp. 303–308. MR MR0098419 (20 #4878)
- [Eli67] Halldór I. Eliasson, *Geometry of manifolds of maps*, J. Differential Geometry **1** (1967), 169–194. MR MR0226681 (37 #2268)
- [Éme89] Michel Émery, *Stochastic calculus in manifolds*, Universitext, Springer-Verlag, Berlin, 1989, With an appendix by P.-A. Meyer. MR MR1030543 (90k:58244)
- [Fey48] R. P. Feynman, *Space-time approach to non-relativistic quantum mechanics*, Rev. Modern Physics **20** (1948), 367–387. MR MR0026940 (10,224b)
- [FK72] P. Flaschel and W. Klingenberg, *Riemannsche Hilbertmannigfaltigkeiten. Periodische Geodätische. Mit einem Anhang von H. Karcher*, Springer-Verlag, Berlin, 1972, Lecture Notes in Mathematics, Vol. 282. MR MR0341527 (49 #6275)
- [Fuj80] Daisuke Fujiwara, *Remarks on convergence of the Feynman path integrals*, Duke Math. J. **47** (1980), no. 3, 559–600. MR MR587166 (83c:81030)
- [Gan64] Ramesh Gangolli, *On the construction of certain diffusions on a differentiable manifold*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete **2** (1964), 406–419 (1964). MR MR0165590 (29 #2870)
- [Gro85] Leonard Gross, *A Poincaré lemma for connection forms*, J. Funct. Anal. **63** (1985), no. 1, 1–46. MR MR795515 (87a:53110)
- [Ich97] Wataru Ichinose, *On the formulation of the Feynman path integral through broken line paths*, Comm. Math. Phys. **189** (1997), no. 1, 17–33. MR MR1478529 (98k:81138)

- [IM85] Atsushi Inoue and Yoshiaki Maeda, *On integral transformations associated with a certain Lagrangian—as a prototype of quantization*, J. Math. Soc. Japan **37** (1985), no. 2, 219–244. MR MR780661 (86h:58054)
- [Kli78] Wilhelm Klingenberg, *Lectures on closed geodesics*, Springer-Verlag, Berlin, 1978, Grundlehren der Mathematischen Wissenschaften, Vol. 230. MR MR0478069 (57 #17563)
- [Kun97] Hiroshi Kunita, *Stochastic flows and stochastic differential equations*, Cambridge Studies in Advanced Mathematics, vol. 24, Cambridge University Press, Cambridge, 1997, Reprint of the 1990 original. MR MR1472487 (98e:60096)
- [Lea97] R. Leandre, *Invariant Sobolev calculus on the free loop space*, Acta Appl. Math. **46** (1997), no. 3, 267–350. MR MR1440476 (98a:60077)
- [NY78] Shintaro Nakao and Yuiti Yamato, *Approximation theorem on stochastic differential equations*, Proceedings of the International Symposium on Stochastic Differential Equations (Res. Inst. Math. Sci., Kyoto Univ., Kyoto, 1976) (New York), Wiley, 1978, pp. 283–296. MR MR536015 (81h:60081)
- [Øks03] Bernt Øksendal, *Stochastic differential equations*, sixth ed., Universitext, Springer-Verlag, Berlin, 2003, An introduction with applications. MR MR2001996 (2004e:60102)
- [Pal63] Richard S. Palais, *Morse theory on Hilbert manifolds*, Topology **2** (1963), 299–340. MR MR0158410 (28 #1633)
- [Pin78] Mark A. Pinsky, *Stochastic Riemannian geometry*, Probabilistic analysis and related topics, Vol. 1, Academic Press, New York, 1978, pp. 199–236. MR MR0501385 (58 #18752)
- [Pro04] Philip E. Protter, *Stochastic integration and differential equations*, second ed., Applications of Mathematics (New York), vol. 21, Springer-Verlag, Berlin, 2004, Stochastic Modelling and Applied Probability. MR MR2020294 (2005k:60008)
- [Rao04] M. M. Rao (ed.), *Real and stochastic analysis*, Trends in Mathematics, Birkhäuser Boston Inc., Boston, MA, 2004, New perspectives. MR MR2090750 (2005d:60002)
- [Res99] Sidney I. Resnick, *A probability path*, Birkhäuser Boston Inc., Boston, MA, 1999. MR MR1664717 (2000b:60002)

- [RS79] Michael Reed and Barry Simon, *Methods of modern mathematical physics. III*, Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1979, Scattering theory. MR MR529429 (80m:81085)
- [RS80] ———, *Methods of modern mathematical physics. I*, second ed., Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1980, Functional analysis. MR MR751959 (85e:46002)
- [Sim79] Barry Simon, *Trace ideals and their applications*, London Mathematical Society Lecture Note Series, vol. 35, Cambridge University Press, Cambridge, 1979. MR MR541149 (80k:47048)
- [Um74] Gregory S. Um, *On normalization problems of the path integral method*, Journal of Mathematical Physics **15** (1974), no. 2, 220–224.
- [WZ65] Eugene Wong and Moshe Zakai, *On the relation between ordinary and stochastic differential equations*, Internat. J. Engrg. Sci. **3** (1965), 213–229. MR MR0183023 (32 #505)
- [WZ67] ———, *On the relation between ordinary and stochastic differential equations and applications to stochastic problems in control theory*, Automatic and remote control III (Proc. Third Congr. Internat. Fed. Automat. Control (IFAC), London, 1966), Vol. 1, p. 5, Paper 3B, Inst. Mech. Engrs., London, 1967, p. 8. MR MR0386015 (52 #6874)

This figure "pos-beg\_\_1.png" is available in "png" format from:

<http://arXiv.org/ps/math/0612711v1>

This figure "pos-beg\_\_2.png" is available in "png" format from:

<http://arXiv.org/ps/math/0612711v1>

This figure "pos-beg\_\_3.png" is available in "png" format from:

<http://arXiv.org/ps/math/0612711v1>